

**Anti-Tuberin TSC2 Antibody Picoband™ (monoclonal, 6I3)**  
Catalog # ABO14893**Specification****Anti-Tuberin TSC2 Antibody Picoband™ (monoclonal, 6I3) - Product Information**

Application	WB, IHC, FC
Primary Accession	<a href="#">P49815</a>
Host	Mouse
Isotype	Mouse IgG2b
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti-Tuberin TSC2 Antibody Picoband™ (monoclonal, 6I3) . Tested in Flow Cytometry, IHC, WB applications. This antibody reacts with Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500 µg/ml.

**Anti-Tuberin TSC2 Antibody Picoband™ (monoclonal, 6I3) - Additional Information**

**Gene ID** 7249

**Other Names**

Tuberin, Tuberous sclerosis 2 protein, TSC2 {ECO:0000303|PubMed:7558029, ECO:0000312|HGNC:HGNC:12363}

**Calculated MW**

280 kDa KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human<br> Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat<br> Flow Cytometry, 1-3 µg/1x10<sup>6</sup> cells, Human<br>

**Subcellular Localization**

Cytoplasm; Membrane; Peripheral membrane protein

**Tissue Specificity**

Liver, brain, heart, lymphocytes, fibroblasts, biliary epithelium, pancreas, skeletal muscle, kidney, lung and placenta.

**Contents**

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

**Immunogen**

E.coli-derived human Tuberin recombinant protein (Position: H1611-V1807). Human Tuberin shares 94% and 90% amino acid (aa) sequence identity with mouse and rat Tuberin, respectively.

### Cross Reactivity

No cross-reactivity with other proteins.

### Storage

Store at  $-20^{\circ}\text{C}$  for one year from date of receipt. After reconstitution, at  $4^{\circ}\text{C}$  for one month. It can also be aliquotted and stored frozen at  $-20^{\circ}\text{C}$  for six months. Avoid repeated freeze-thaw cycles.

## Anti-Tuberin TSC2 Antibody Picoband™ (monoclonal, 6I3) - Protein Information

**Name** TSC2 {ECO:0000303|PubMed:7558029, ECO:0000312|HGNC:HGNC:12363}

### Function

Catalytic component of the TSC-TBC complex, a multiprotein complex that acts as a negative regulator of the canonical mTORC1 complex, an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:<a href="http://www.uniprot.org/citations/12172553" target="\_blank">12172553</a>, PubMed:<a href="http://www.uniprot.org/citations/12271141" target="\_blank">12271141</a>, PubMed:<a href="http://www.uniprot.org/citations/12842888" target="\_blank">12842888</a>, PubMed:<a href="http://www.uniprot.org/citations/12906785" target="\_blank">12906785</a>, PubMed:<a href="http://www.uniprot.org/citations/15340059" target="\_blank">15340059</a>, PubMed:<a href="http://www.uniprot.org/citations/22819219" target="\_blank">22819219</a>, PubMed:<a href="http://www.uniprot.org/citations/24529379" target="\_blank">24529379</a>, PubMed:<a href="http://www.uniprot.org/citations/28215400" target="\_blank">28215400</a>, PubMed:<a href="http://www.uniprot.org/citations/33436626" target="\_blank">33436626</a>, PubMed:<a href="http://www.uniprot.org/citations/35772404" target="\_blank">35772404</a>). Within the TSC-TBC complex, TSC2 acts as a GTPase-activating protein (GAP) for the small GTPase RHEB, a direct activator of the protein kinase activity of mTORC1 (PubMed:<a href="http://www.uniprot.org/citations/12172553" target="\_blank">12172553</a>, PubMed:<a href="http://www.uniprot.org/citations/12820960" target="\_blank">12820960</a>, PubMed:<a href="http://www.uniprot.org/citations/12842888" target="\_blank">12842888</a>, PubMed:<a href="http://www.uniprot.org/citations/12906785" target="\_blank">12906785</a>, PubMed:<a href="http://www.uniprot.org/citations/15340059" target="\_blank">15340059</a>, PubMed:<a href="http://www.uniprot.org/citations/22819219" target="\_blank">22819219</a>, PubMed:<a href="http://www.uniprot.org/citations/24529379" target="\_blank">24529379</a>, PubMed:<a href="http://www.uniprot.org/citations/33436626" target="\_blank">33436626</a>). In absence of nutrients, the TSC-TBC complex inhibits mTORC1, thereby preventing phosphorylation of ribosomal protein S6 kinase (RPS6KB1 and RPS6KB2) and EIF4EBP1 (4E-BP1) by the mTORC1 signaling (PubMed:<a href="http://www.uniprot.org/citations/12172553" target="\_blank">12172553</a>, PubMed:<a href="http://www.uniprot.org/citations/12271141" target="\_blank">12271141</a>, PubMed:<a href="http://www.uniprot.org/citations/12842888" target="\_blank">12842888</a>, PubMed:<a href="http://www.uniprot.org/citations/12906785" target="\_blank">12906785</a>, PubMed:<a href="http://www.uniprot.org/citations/22819219" target="\_blank">22819219</a>, PubMed:<a href="http://www.uniprot.org/citations/24529379" target="\_blank">24529379</a>, PubMed:<a href="http://www.uniprot.org/citations/28215400" target="\_blank">28215400</a>, PubMed:<a href="http://www.uniprot.org/citations/35772404" target="\_blank">35772404</a>). The TSC-TBC complex is inactivated in response to nutrients, relieving inhibition of mTORC1 (PubMed:<a href="http://www.uniprot.org/citations/12172553" target="\_blank">12172553</a>, PubMed:<a href="http://www.uniprot.org/citations/24529379" target="\_blank">24529379</a>). Involved in microtubule-mediated protein transport via its ability to regulate mTORC1 signaling (By similarity). Also stimulates the intrinsic GTPase activity of the Ras- related proteins RAP1A and RAB5 (By similarity).

### Cellular Location

Lysosome membrane; Peripheral membrane protein. Cytoplasm, cytosol Note=Recruited to lysosomal membranes in a RHEB-dependent process in absence of nutrients (PubMed:24529379). In response to insulin signaling and phosphorylation by PKB/AKT1, the complex dissociates from lysosomal membranes and relocates to the cytosol (PubMed:24529379)

#### Tissue Location

Liver, brain, heart, lymphocytes, fibroblasts, biliary epithelium, pancreas, skeletal muscle, kidney, lung and placenta.

### Anti-Tuberin TSC2 Antibody Picoband™ (monoclonal, 6I3) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-Tuberin TSC2 Antibody Picoband™ (monoclonal, 6I3) - Images




Figure 1. Western blot analysis of TSC2 using anti-TSC2 antibody (M00229-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human HEK293 tissue lysates,

Lane 2: human PC-3 whole cell lysates,

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-TSC2 antigen affinity purified polyclonal antibody (Catalog # M00229-1) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for

TSC2 at approximately 280KD. The expected band size for TSC2 is at 201KD.




Figure 2. IHC analysis of TSC2 using anti-TSC2 antibody (M00229-1). TSC2 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu\text{g}/\text{ml}$  mouse anti-TSC2 Antibody (M00229-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit (Catalog # SV0001) with DAB as the chromogen.




Figure 3. Flow Cytometry analysis of A549 cells using anti-TSC2 antibody (M00229-1). Overlay histogram showing A549 cells stained with M00229-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-TSC2 Antibody (M00229-1, 1  $\mu\text{g}/1 \times 10^6$  cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10  $\mu\text{g}/1 \times 10^6$  cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1  $\mu\text{g}/1 \times 10^6$ ) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

### Anti-Tuberin TSC2 Antibody Picoband™ (monoclonal, 6I3) - Background

Tuberous sclerosis 2 protein, also known as TSC2 or Tuberin is a protein that is in humans. The chromosomal location of this gene is 16p13.3. Mutations in this gene lead to tuberous sclerosis complex. Its gene product is believed to be a tumor suppressor and is able to stimulate specific GTPases. The protein associates with hamartin in a cytosolic complex, possibly acting as a chaperone for hamartin. This gene involved in microtubule-mediated protein transport, but this seems to be due to unregulated mTOR signaling. It stimulates weakly the intrinsic GTPase activity of the Ras-related proteins RAP1A and RAB5 in vitro.