

Anti-CD5 Antibody Picoband™ (monoclonal, 4E2)
Catalog # ABO14898

Specification

Anti-CD5 Antibody Picoband™ (monoclonal, 4E2) - Product Information

Application	WB, IHC
Primary Accession	P06127
Host	Mouse
Isotype	Mouse IgG1
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-CD5 Antibody Picoband™ (monoclonal, 4E2) . Tested in IHC, WB applications. This antibody reacts with Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500 µg/ml.

Anti-CD5 Antibody Picoband™ (monoclonal, 4E2) - Additional Information

Gene ID 921

Other Names

T-cell surface glycoprotein CD5, Lymphocyte antigen T1/Leu-1, CD5, CD5, LEU1

Calculated MW

67 kDa KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human
 Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat

Subcellular Localization

Cell membrane; Single-pass type I membrane protein

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E. coli-derived human CD5 recombinant protein (Position: R25-L495).

Cross Reactivity

No cross-reactivity with other proteins.

Storage

Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored

frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.

Anti-CD5 Antibody Picoband™ (monoclonal, 4E2) - Protein Information

Name CD5

Synonyms LEU1

Function

Lymphoid-specific receptor expressed by all T-cells and in a subset of B-cells known as B1a cells. Plays a role in the regulation of TCR and BCR signaling, thymocyte selection, T-cell effector differentiation and immune tolerance. Acts by interacting with several ligands expressed on B-cells such as CD5L or CD72 and thereby plays an important role in contact-mediated, T-dependent B-cell activation and in the maintenance of regulatory T and B-cell homeostasis. Functions as a negative regulator of TCR signaling during thymocyte development by associating with several signaling proteins including LCK, CD3Z chain, PI3K or CBL (PubMed:1385158, PubMed:1384049). Mechanistically, co- engagement of CD3 with CD5 enhances phosphorylated CBL recruitment leading to increased VAV1 phosphorylation and degradation (PubMed:23376399). Modulates B-cell biology through ERK1/2 activation in a Ca(2+)-dependent pathway via the non-selective Ca(2+) channel TRPC1, leading to IL-10 production (PubMed:27499044).

Cellular Location


Cell membrane {ECO:0000250|UniProtKB:P13379}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P13379}

Anti-CD5 Antibody Picoband™ (monoclonal, 4E2) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD5 Antibody Picoband™ (monoclonal, 4E2) - Images






Figure 1. Western blot analysis of CD5 using anti-CD5 antibody (M00480-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human Jurkat tissue lysates,

Lane 2: human CCRM-CEM whole cell lysates,

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-CD5 antigen affinity purified polyclonal antibody (Catalog # M00480-2) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for CD5 at approximately 67kD. The expected band size for CD5 is at 54kD.




Figure 2. IHC analysis of CD5 using anti-CD5 antibody (M00480-2).

CD5 was detected in paraffin-embedded section of human tonsil cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 µg/ml mouse anti-CD5 Antibody (M00480-2) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.

Anti-CD5 Antibody Picoband™ (monoclonal, 4E2) - Background

CD5 is a member of the scavenger receptor cysteine-rich (SRCR) superfamily. Members of this family are secreted or membrane-anchored proteins mainly found in cells associated with the

immune system. In humans, the gene is located on the long arm of chromosome 11. This protein is a type-I transmembrane glycoprotein found on the surface of thymocytes, T lymphocytes and a subset of B lymphocytes. The encoded protein contains three SRCR domains and may act as a receptor to regulate T-cell proliferation. Alternative splicing results in multiple transcript variants encoding different isoforms.