

**Anti-PROM1/CD133 Antibody Picoband™ (monoclonal, 7F13)**  
Catalog # ABO14992**Specification****Anti-PROM1/CD133 Antibody Picoband™ (monoclonal, 7F13) - Product Information**

Application	WB
Primary Accession	<a href="#">O43490</a>
Host	Mouse
Isotype	Mouse IgG1
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti-PROM1/CD133 Antibody Picoband™ (monoclonal, 7F13) . Tested in WB applications. This antibody reacts with Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-PROM1/CD133 Antibody Picoband™ (monoclonal, 7F13) - Additional Information**

**Gene ID** 8842

**Other Names**

Prominin-1, Antigen AC133, Prominin-like protein 1, CD133, PROM1, PROML1

**Calculated MW**

120 kDa KDa

**Application Details**

Western blot, 0.25-0.5 µg/ml, Human<br>

**Contents**

Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na<sub>2</sub>HPO<sub>4</sub>.

**Immunogen**

E.coli-derived human PROM1/CD133 recombinant protein (Position: P531-H865).  
HumanPROM1/CD133 shares 61% amino acid (aa) sequence identity with mouse PROM1/CD133.

**Purification**

Immunogen affinity purified.

**Storage**

**Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.**

**Anti-PROM1/CD133 Antibody Picoband™ (monoclonal, 7F13) - Protein Information**

**Name** PROM1

**Synonyms** PROML1

**Function**

May play a role in cell differentiation, proliferation and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/24556617" target="\_blank">24556617</a>). Binds cholesterol in cholesterol- containing plasma membrane microdomains and may play a role in the organization of the apical plasma membrane in epithelial cells. During early retinal development acts as a key regulator of disk morphogenesis. Involved in regulation of MAPK and Akt signaling pathways. In neuroblastoma cells suppresses cell differentiation such as neurite outgrowth in a RET-dependent manner (PubMed:<a href="http://www.uniprot.org/citations/20818439" target="\_blank">20818439</a>).

**Cellular Location**

Apical cell membrane; Multi-pass membrane protein. Cell projection, microvillus membrane; Multi-pass membrane protein. Cell projection, cilium, photoreceptor outer segment Endoplasmic reticulum. Endoplasmic reticulum-Golgi intermediate compartment. Note=Found in extracellular membrane particles in various body fluids such as cerebrospinal fluid, saliva, seminal fluid and urine

**Tissue Location**


Isoform 1 is selectively expressed on CD34 hematopoietic stem and progenitor cells in adult and fetal bone marrow, fetal liver, cord blood and adult peripheral blood. Isoform 1 is not detected on other blood cells. Isoform 1 is also expressed in a number of non-lymphoid tissues including retina, pancreas, placenta, kidney, liver, lung, brain and heart. Found in saliva within small membrane particles. Isoform 2 is predominantly expressed in fetal liver, skeletal muscle, kidney, and heart as well as adult pancreas, kidney, liver, lung, and placenta. Isoform 2 is highly expressed in fetal liver, low in bone marrow, and barely detectable in peripheral blood Isoform 2 is expressed on hematopoietic stem cells and in epidermal basal cells (at protein level). Expressed in adult retina by rod and cone photoreceptor cells (at protein level)

**Anti-PROM1/CD133 Antibody Picoband™ (monoclonal, 7F13) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-PROM1/CD133 Antibody Picoband™ (monoclonal, 7F13) - Images**






Figure 1. Western blot analysis of PROM1/CD133 using anti-PROM1/CD133 antibody (M01767-4). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human CACO-2 whole cell lysates,  
 Lane 2: human SW620 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-PROM1/CD133 antigen affinity purified monoclonal antibody (Catalog # M01767-4) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for PROM1/CD133 at approximately 120KD. The expected band size for PROM1/CD133 is at 120KD.

**Anti-PROM1/CD133 Antibody Picoband™ (monoclonal, 7F13) - Background**

Prominin-1, also known as CD133, is a glycoprotein that in humans is encoded by the PROM1 gene. It is mapped to 4p15.32. Prominin-1 is a member of pentaspan transmembrane glycoproteins (5-transmembrane, 5-TM), which specifically localize to cellular protrusions. This gene encodes a pentaspan transmembrane glycoprotein. The protein localizes to membrane protrusions and is often expressed on adult stem cells, where it is thought to function in maintaining stem cell properties by suppressing differentiation. It has been proposed to act as an organizer of cell membrane topology. Prominin-1 was expressed not only on metastatic colon cancer cells, but also on differentiated colonic epithelium in both adult mice and humans.