

Anti-Transketolase/TKT Picoband[™] Antibody (monoclonal, 213)

Catalog # ABO15006

Specification

Anti-Transketolase/TKT Picoband[™] Antibody (monoclonal, 213) - Product Information

Application Primary Accession Host Isotype Reactivity Clonality Format **Description** Anti-Transketolase/TKT Pi WB, IHC, IF, ICC, FC <u>P29401</u> Mouse Mouse IgG2b Rat, Human, Mouse Monoclonal Lyophilized

Anti-Transketolase/TKT Picoband[™] Antibody (monoclonal, 213) . Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Transketolase/TKT Picoband[™] Antibody (monoclonal, 213) - Additional Information

Gene ID 7086

Other Names Transketolase, TK, 2.2.1.1, TKT

Calculated MW 68 kDa KDa

Application Details Western blot, 0.1-0.25 μg/ml, Human, Mouse, Rat
 Immunohistochemistry (Paraffin-embedded Section), 1-2 μg/ml, Human
 Immunocytochemistry/Immunofluorescence, 5 μg/ml, Human
 Flow Cytometry, 1-3 μg/1x10^6 cells, Human

Contents Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na2HPO4.

Immunogen E.coli-derived human Transketolase/TKT recombinant protein (Position: M1-A116).

Purification Immunogen affinity purified.

Storage

Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.



Anti-Transketolase/TKT Picoband[™] Antibody (monoclonal, 213) - Protein Information

Name TKT

Function

Catalyzes the transfer of a two-carbon ketol group from a ketose donor to an aldose acceptor, via a covalent intermediate with the cofactor thiamine pyrophosphate.

Anti-Transketolase/TKT Picoband[™] Antibody (monoclonal, 213) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Transketolase/TKT Picoband[™] Antibody (monoclonal, 2I3) - Images



Figure 1. Western blot analysis of Transketolase/TKT using anti-Transketolase/TKT antibody (M02197-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human A549 whole cell lysates,

Lane 2: human SW620 whole cell lysates,

Lane 3: human Raji whole cell lysates,

Lane 4: rat liver tissue lysates,

Lane 5: rat RH35 whole cell lysates,

Lane 6: mouse liver tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Transketolase/TKT antigen affinity purified monoclonal antibody (Catalog # M02197-1) at 0.25 μ g/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for



Transketolase/TKT at approximately 68KD. The expected band size for Transketolase/TKT is at 68KD.



Figure 2. IHC analysis of Transketolase/TKT using anti-Transketolase/TKT antibody (M02197-1). Transketolase/TKT was detected in paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-Transketolase/TKT Antibody (M02197-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.



Figure 3. IHC analysis of Transketolase/TKT using anti-Transketolase/TKT antibody (M02197-1). Transketolase/TKT was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-Transketolase/TKT Antibody (M02197-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.





Figure 4. IHC analysis of Transketolase/TKT using anti-Transketolase/TKT antibody (M02197-1). Transketolase/TKT was detected in paraffin-embedded section of human pancreatic cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-Transketolase/TKT Antibody (M02197-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.



Figure 5. IHC analysis of Transketolase/TKT using anti-Transketolase/TKT antibody (M02197-1). Transketolase/TKT was detected in paraffin-embedded section of human thyroid cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-Transketolase/TKT Antibody (M02197-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.



Figure 6. IF analysis of Transketolase/TKT using anti-Transketolase/TKT antibody (M02197-1). Transketolase/TKT was detected in immunocytochemical section of A549 cells. Enzyme antigen



retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 μ g/mL mouse anti-Transketolase/TKT Antibody (M02197-1) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



Figure 7. Flow Cytometry analysis of THP-1 cells using anti-Transketolase/TKT antibody (M02197-1).

Overlay histogram showing THP-1 cells stained with M02197-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-Transketolase/TKT Antibody (M02197-1, 1 μ g/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 μ g/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 μ g/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



Figure 8. Flow Cytometry analysis of A549 cells using anti-Transketolase/TKT antibody (M02197-1).

Overlay histogram showing A549 cells stained with M02197-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-Transketolase/TKT Antibody (M02197-1, 1 μ g/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 μ g/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 μ g/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Anti-Transketolase/TKT Picoband[™] Antibody (monoclonal, 213) - Background

Transketolase is a thiamine-dependent enzyme that links the pentose phosphate pathway with the glycolytic pathway. The pentose phosphate pathway, which is active in most tissues, provides sugar phosphates for intermediary biosynthesis, especially nucleotide metabolism, and generates the biosynthetic reducing power for the cell in the form of NADPH. Transketolase is directly involved in the branch of the pathway that channels excess sugar phosphates to glycolysis, enabling the



production of NADPH to be maintained under different metabolic conditions. NADPH is critical for maintaining cerebral glutathione, and thus it is likely that transketolase plays an important role in brain metabolism.