

**Anti- IL-9 Monoclonal Antibody**  
Catalog # ABO15022**Specification**

---

**Anti- IL-9 Monoclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P15248</a>
Host	Mouse
Isotype	Mouse IgG2b, $\kappa$
Reactivity	Rat, Mouse
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti- IL-9 Monoclonal Antibody . Tested in WB applications. This antibody reacts with Mouse, Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500  $\mu$ g/ml.

**Anti- IL-9 Monoclonal Antibody - Additional Information**

**Gene ID** 3578

**Other Names**

Interleukin-9, IL-9, Cytokine P40, T-cell growth factor P40, IL9

**Calculated MW**

16 kDa KDa

**Application Details**

Western blot, 0.25-0.5 $\mu$ g/ml, Mouse, Rat

**Protein Name**

Interleukin-9

**Contents**

PBS, pH 7.0. Contains no stabilizers or preservatives

**Immunogen**

Human IL-9 coupled to ovalbumin

**Purification**

Immunogen affinity purified.

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti- IL-9 Monoclonal Antibody - Protein Information**

---

## Name IL9

### Function

Multifunctional cytokine secreted mainly by T-helper 2 lymphocytes and also mast cells or NKT cells that plays important roles in the immune response against parasites (PubMed:<a href="http://www.uniprot.org/citations/29742432" target="\_blank">29742432</a>). Affects intestinal epithelial permeability and adaptive immunity (PubMed:<a href="http://www.uniprot.org/citations/29742432" target="\_blank">29742432</a>). In addition, induces the differentiation of specific T-cell subsets such as IL-17 producing helper T-cells (TH17) and also proliferation and differentiation of mast cells. Mechanistically, exerts its biological effects through a receptor composed of IL9R subunit and a signal transducing subunit IL2RG. Receptor stimulation results in the rapid activation of JAK1 and JAK3 kinase activities leading to STAT1, STAT3 and STAT5-mediated transcriptional programs. Induction of differentiation genes seems to be mediated by STAT1 alone, while protection of cells from apoptosis depends on STAT3 and STAT5.

### Cellular Location

Secreted.

## Anti- IL-9 Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti- IL-9 Monoclonal Antibody - Images




Figure 1. Western blot analysis of IL-9 using anti-IL-9 antibody (M02925).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: rat spleen tissue lysates,

Lane 2: rat thymus tissue lysates,  
Lane 3: mouse thymus tissue lysates,  
Lane 4: mouse RAW264.7 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-IL-9 antigen affinity purified monoclonal antibody (Catalog # M02925) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for IL-9 at approximately 16KD. The expected band size for IL-9 is at 16KD.

#### **Anti- IL-9 Monoclonal Antibody - Background**

The protein encoded by this gene is a cytokine that acts as a regulator of a variety of hematopoietic cells. This cytokine stimulates cell proliferation and prevents apoptosis. It functions through the interleukin 9 receptor (IL9R), which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes. The gene encoding this cytokine has been identified as a candidate gene for asthma. Genetic studies on a mouse model of asthma demonstrated that this cytokine is a determining factor in the pathogenesis of bronchial hyperresponsiveness. [provided by RefSeq, Jul 2008]