

**Anti- IFN Gamma Monoclonal Antibody**  
**Catalog # ABO15025****Specification**

---

**Anti- IFN Gamma Monoclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P01579</a>
Host	Mouse
Isotype	Mouse IgG1
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti- IFN Gamma Monoclonal Antibody . Tested in WB applications. This antibody reacts with Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500 µg/ml.

**Anti- IFN Gamma Monoclonal Antibody - Additional Information**

**Gene ID** 3458

**Other Names**

Interferon gamma, IFN-gamma, Immune interferon, IFNG

**Application Details**

Western blot, 0.25-0.5 µg/ml, Human

**Protein Name**

interferon gamma

**Contents**

PBS, pH 7.0. Contains no stabilizers or preservatives

**Immunogen**

Recombinant human IFN $\gamma$

**Purification**

Immunogen affinity purified.

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti- IFN Gamma Monoclonal Antibody - Protein Information**

**Name IFNG****Function**

Type II interferon produced by immune cells such as T-cells and NK cells that plays crucial roles in antimicrobial, antiviral, and antitumor responses by activating effector immune cells and enhancing antigen presentation (PubMed:<a href="http://www.uniprot.org/citations/16914093" target="\_blank">16914093</a>, PubMed:<a href="http://www.uniprot.org/citations/8666937" target="\_blank">8666937</a>). Primarily signals through the JAK-STAT pathway after interaction with its receptor IFNGR1 to affect gene regulation (PubMed:<a href="http://www.uniprot.org/citations/8349687" target="\_blank">8349687</a>). Upon IFNG binding, IFNGR1 intracellular domain opens out to allow association of downstream signaling components JAK2, JAK1 and STAT1, leading to STAT1 activation, nuclear translocation and transcription of IFNG-regulated genes. Many of the induced genes are transcription factors such as IRF1 that are able to further drive regulation of a next wave of transcription (PubMed:<a href="http://www.uniprot.org/citations/16914093" target="\_blank">16914093</a>). Plays a role in class I antigen presentation pathway by inducing a replacement of catalytic proteasome subunits with immunoproteasome subunits (PubMed:<a href="http://www.uniprot.org/citations/8666937" target="\_blank">8666937</a>). In turn, increases the quantity, quality, and repertoire of peptides for class I MHC loading (PubMed:<a href="http://www.uniprot.org/citations/8163024" target="\_blank">8163024</a>). Increases the efficiency of peptide generation also by inducing the expression of activator PA28 that associates with the proteasome and alters its proteolytic cleavage preference (PubMed:<a href="http://www.uniprot.org/citations/11112687" target="\_blank">11112687</a>). Up-regulates as well MHC II complexes on the cell surface by promoting expression of several key molecules such as cathepsins B/CTSB, H/CTSH, and L/CTSL (PubMed:<a href="http://www.uniprot.org/citations/7729559" target="\_blank">7729559</a>). Participates in the regulation of hematopoietic stem cells during development and under homeostatic conditions by affecting their development, quiescence, and differentiation (By similarity).

**Cellular Location**

Secreted.


**Tissue Location**

Released primarily from activated T lymphocytes.

**Anti- IFN Gamma Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti- IFN Gamma Monoclonal Antibody - Images**




Figure 1. Western blot analysis of IFN Gamma using anti-IFN Gamma antibody (M00393-4). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours.

Lane 1: recombinant human IFN Gamma protein 15ng,  
Lane 2: recombinant human IFN Gamma protein 10ng.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-IFN Gamma antigen affinity purified monoclonal antibody (Catalog # M00393-4) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system.

#### **Anti- IFN Gamma Monoclonal Antibody - Background**

Interferon-gamma (IFN-gamma) is an inflammatory cytokine that has been implicated in the development of fibrosis in inflamed tissues. The production of IFN-gamma, which is under genetic control, can influence the development of fibrosis in lung allografts. IFN-gamma is also produced by natural killer (NK) cells and most prominently by CD8 cytotoxic T cells, and is vital for the control of microbial pathogens. Interferon gamma is believed to be crucial for host defence against many infections. Genetically determined variability in IFN-gamma and expression might be important for the development of tuberculosis. IFN-gamma activates human macrophage oxidative metabolism and antimicrobial activity. In addition to having antiviral activity, IFN-gamma has important immunoregulatory functions. IFN-gamma plays an important role in the control of neointima proliferation.