

**Anti-PNP Antibody Picoband™ (monoclonal, 7G5)**  
**Catalog # ABO15037****Specification**

---

**Anti-PNP Antibody Picoband™ (monoclonal, 7G5) - Product Information**

Application	WB, IHC, IF, ICC
Primary Accession	<a href="#">P00491</a>
Host	Mouse
Isotype	Mouse IgG2b
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti-PNP Antibody Picoband™ (monoclonal, 7G5) . Tested in IF, IHC, ICC, WB applications. This antibody reacts with Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-PNP Antibody Picoband™ (monoclonal, 7G5) - Additional Information**

**Gene ID** 4860

**Other Names**

Purine nucleoside phosphorylase, PNP, 2.4.2.1, Inosine phosphorylase, Inosine-guanosine phosphorylase, PNP, NP

**Calculated MW**

32 kDa KDa

**Application Details**

Western blot, 0.25-0.5 µg/ml, Human<br> Immunohistochemistry (Paraffin-embedded Section), 2-5 µg/ml, Human<br> Immunocytochemistry/Immunofluorescence, 5 µg/ml, Human<br>

**Contents**

Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na<sub>2</sub>HPO<sub>4</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence in the middle region of human PNP, different from the related mouse sequence by six amino acids, and from the related rat sequence by five amino acids.

**Purification**

Immunogen affinity purified.

**Storage**

**Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid**

repeated freeze-thaw cycles.

## **Anti-PNP Antibody Picoband™ (monoclonal, 7G5) - Protein Information**

**Name** PNP

**Synonyms** NP

### **Function**

Catalyzes the phosphorolytic breakdown of the N-glycosidic bond in the beta-(deoxy)ribonucleoside molecules, with the formation of the corresponding free purine bases and pentose-1-phosphate (PubMed:<a href="http://www.uniprot.org/citations/23438750" target="\_blank">23438750</a>, PubMed:<a href="http://www.uniprot.org/citations/9305964" target="\_blank">9305964</a>). Preferentially acts on 6-oxopurine nucleosides including inosine and guanosine (PubMed:<a href="http://www.uniprot.org/citations/9305964" target="\_blank">9305964</a>).

### **Cellular Location**

Cytoplasm.

### **Tissue Location**


Expressed in red blood cells; overexpressed in red blood cells (cytoplasm) of patients with hereditary non-spherocytic hemolytic anemia of unknown etiology.

## **Anti-PNP Antibody Picoband™ (monoclonal, 7G5) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-PNP Antibody Picoband™ (monoclonal, 7G5) - Images**






Figure 1. Western blot analysis of PNP using anti-PNP antibody (M00988).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30ug of sample under reducing conditions.

Lane 1: human HEK293 whole cell lysates,

Lane 2: human PC-3 whole cell lysates,

Lane 3: human U87 whole cell lysates,

Lane 4: human A549 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-PNP antigen affinity purified monoclonal antibody (Catalog # M00988) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for PNP at approximately 32KD. The expected band size for PNP is at 32KD.




Figure 2. IHC analysis of PNP using anti-PNP antibody (M00988).

PNP was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-PNP Antibody (M00988) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 3. IHC analysis of PNP using anti-PNP antibody (M00988).

PNP was detected in paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-PNP Antibody (M00988) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 4. IF analysis of PNP using anti-PNP antibody (M00988).

PNP was detected in immunocytochemical section of A431 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 µg/mL mouse anti-PNP Antibody (M00988) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

#### **Anti-PNP Antibody Picoband™ (monoclonal, 7G5) - Background**

The PNP gene encodes purine nucleoside phosphorylase, an enzyme that catalyzes the reversible phosphorolysis of the purine nucleosides and deoxynucleosides inosine, guanosine, deoxyinosine, and deoxyguanosine. It is presented results from gene dosage studies consistent with assignment of the PNP locus to band 14q13. PNP is expressed in most tissues, with markedly greater expression in lymphoid tissues. Genetic deficiencies of PNP result in severely compromised Tlymphocyte function and neurologic dysfunction.