

Anti-Cytokeratin 5 Antibody Picoband™ (monoclonal, 9H2F8)
Catalog # ABO15095**Specification****Anti-Cytokeratin 5 Antibody Picoband™ (monoclonal, 9H2F8) - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	P13647
Host	Mouse
Isotype	Mouse IgG2a
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-Cytokeratin 5 Antibody Picoband™ (monoclonal, 9H2F8) . Tested in FCM, IF, IHC, ICC, WB applications. This antibody reacts with Human.

Reconstitution

Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

Anti-Cytokeratin 5 Antibody Picoband™ (monoclonal, 9H2F8) - Additional Information

Gene ID 3852

Other Names

Keratin, type II cytoskeletal 5, 58 kDa cyokeratin, Cytokeratin-5, CK-5, Keratin-5, K5, Type-II keratin Kb5, KRT5

Calculated MW

62 kDa KDa

Application Details

Western blot, 0.25-0.5 µg/ml, Human
 Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/ml, Human
 Immunocytochemistry/Immunofluorescence, 5 µg/ml, Human
 Flow Cytometry, 1-3 µg/1x⁶ cells, Human

Contents

Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na₂HPO₄.

Immunogen

A synthetic peptide corresponding to a sequence in the middle region of human Cytokeratin 5, different from the related mouse sequence by one amino acid, and identical to the related rat sequence.

Purification

Immunogen affinity purified.

Storage

**At -20°C for one year from date of receipt.
After reconstitution, at 4°C for one month.
It can also be aliquotted and stored frozen**

at -20°C for six months. Avoid repeated freezing and thawing.

Anti-Cytokeratin 5 Antibody Picoband™ (monoclonal, 9H2F8) - Protein Information

Name KRT5

Function

Required for the formation of keratin intermediate filaments in the basal epidermis and maintenance of the skin barrier in response to mechanical stress (By similarity). Regulates the recruitment of Langerhans cells to the epidermis, potentially by modulation of the abundance of macrophage chemotactic cytokines, macrophage inflammatory cytokines and CTNND1 localization in keratinocytes (By similarity).

Cellular Location

Cytoplasm.

Tissue Location

Expressed in corneal epithelium (at protein level) (PubMed:26758872). Expressed in keratinocytes (at protein level) (PubMed:20128788, PubMed:31302245).

Anti-Cytokeratin 5 Antibody Picoband™ (monoclonal, 9H2F8) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Cytokeratin 5 Antibody Picoband™ (monoclonal, 9H2F8) - Images




Figure 1. Western blot analysis of Cytokeratin 5 using anti-Cytokeratin 5 antibody (M00398-7). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human A431 whole cell lysates,
Lane 2: human SH-SY5Y whole cell lysates,
Lane 3: human Hacat whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Cytokeratin 5 antigen affinity purified monoclonal antibody (Catalog # M00398-7) at 0.5 μ g/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for Cytokeratin 5 at approximately 62 kDa. The expected band size for Cytokeratin 5 is at 62 kDa.




Figure 2. IHC analysis of Cytokeratin 5 using anti-Cytokeratin 5 antibody (M00398-7). Cytokeratin 5 was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-Cytokeratin 5 Antibody (M00398-7) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 3. IHC analysis of Cytokeratin 5 using anti-Cytokeratin 5 antibody (M00398-7). Cytokeratin 5 was detected in a paraffin-embedded section of human metaplasia of squamous cells of the renal pelvis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-Cytokeratin 5 Antibody (M00398-7) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30

minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 4. IHC analysis of Cytokeratin 5 using anti-Cytokeratin 5 antibody (M00398-7). Cytokeratin 5 was detected in a paraffin-embedded section of human rectal moderately differentiated adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Cytokeratin 5 Antibody (M00398-7) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 5. IF analysis of Cytokeratin 5 using anti-Cytokeratin 5 antibody (M00398-7). Cytokeratin 5 was detected in an immunocytochemical section of A431 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 µg/mL mouse anti-Cytokeratin 5 Antibody (M00398-7) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 6. Flow Cytometry analysis of A431 cells using anti-Cytokeratin 5 antibody (M00398-7). Overlay histogram showing A431 cells stained with M00398-7 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-Cytokeratin 5 Antibody (M00398-7, 1 $\mu\text{g}/1 \times 10^6$ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 $\mu\text{g}/1 \times 10^6$ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 $\mu\text{g}/1 \times 10^6$) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Anti-Cytokeratin 5 Antibody Picoband™ (monoclonal, 9H2F8) - Background

Cytokeratin 5, also known as KRT5, K5, or CK5, is a protein that is encoded in humans by the KRT5 gene. The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the basal layer of the epidermis with family member KRT14. Mutations in these genes have been associated with a complex of diseases termed epidermolysis bullosa simplex. The type II cytokeratins are clustered in a region of chromosome 12q12-q13.