

**Anti-PI3 Kinase p85 alpha Rabbit Monoclonal Antibody**  
Catalog # ABO15162**Specification****Anti-PI3 Kinase p85 alpha Rabbit Monoclonal Antibody - Product Information**

Application	WB, IF, ICC
Primary Accession	<a href="#">P27986</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-PI3 Kinase p85 alpha Rabbit Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

**Anti-PI3 Kinase p85 alpha Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 5295

**Other Names**

Phosphatidylinositol 3-kinase regulatory subunit alpha, PI3-kinase regulatory subunit alpha, PI3K regulatory subunit alpha, PtdIns-3-kinase regulatory subunit alpha, Phosphatidylinositol 3-kinase 85 kDa regulatory subunit alpha, PI3-kinase subunit p85-alpha, PtdIns-3-kinase regulatory subunit p85-alpha, PIK3R1, GRB1

**Calculated MW**

84 kDa KDa

**Application Details**

WB 1:500-1:2000<br>ICC/IF 1:50-1:200

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human PI3 Kinase p85 alpha

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-PI3 Kinase p85 alpha Rabbit Monoclonal Antibody - Protein Information**

**Name** PIK3R1

**Synonyms** GRB1

**Function**

Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Plays an important role in signaling in response to FGFR1, FGFR2, FGFR3, FGFR4, KITLG/SCF, KIT, PDGFRA and PDGFRB. Likewise, plays a role in ITGB2 signaling (PubMed:<a href="http://www.uniprot.org/citations/17626883" target="\_blank">17626883</a>, PubMed:<a href="http://www.uniprot.org/citations/19805105" target="\_blank">19805105</a>, PubMed:<a href="http://www.uniprot.org/citations/7518429" target="\_blank">7518429</a>). Modulates the cellular response to ER stress by promoting nuclear translocation of XBP1 isoform 2 in a ER stress- and/or insulin-dependent manner during metabolic overloading in the liver and hence plays a role in glucose tolerance improvement (PubMed:<a href="http://www.uniprot.org/citations/20348923" target="\_blank">20348923</a>).

**Tissue Location**

Isoform 2 is expressed in skeletal muscle and brain, and at lower levels in kidney and cardiac muscle. Isoform 2 and isoform 4 are present in skeletal muscle (at protein level)

**Anti-PI3 Kinase p85 alpha Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-PI3 Kinase p85 alpha Rabbit Monoclonal Antibody - Images**




Figure 1. Western blot analysis of PI3 Kinase p85 alpha using anti-PI3 Kinase p85 alpha antibody (M00318-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing

conditions.

Lane 1: human Jurkat whole cell lysates,

Lane 2: human K562 whole cell lysates,

Lane 3: human 293T whole cell lysates,


Lane 4: rat brain tissue lysates,

Lane 5: rat PC-12 whole cell lysates,


Lane 6: mouse brain tissue lysates,

Lane 7: mouse RAW264.7 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PI3 Kinase p85 alpha antigen affinity purified monoclonal antibody (Catalog # M00318-2) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PI3 Kinase p85 alpha at approximately 84 kDa. The expected band size for PI3 Kinase p85 alpha is at 84 kDa.



Immunofluorescent analysis using the Antibody at 1:50 dilution.



Immunofluorescent analysis using the Antibody at 1:150 dilution.