

### **Anti-MMP17 Rabbit Monoclonal Antibody**

**Catalog # ABO15252** 

### **Specification**

# **Anti-MMP17 Rabbit Monoclonal Antibody - Product Information**

Application WB, IHC, IF, ICC

Primary Accession
Host
Rabbit
Isotype
IgG

Reactivity Rat, Human, Mouse

Clonality Monoclonal Format Liquid

**Description** 

Anti-MMP17 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

# **Anti-MMP17 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID 4326** 

#### **Other Names**

Matrix metalloproteinase-17, MMP-17, 3.4.24.-, Membrane-type matrix metalloproteinase 4, MT-MMP 4, MTMMP4, Membrane-type-4 matrix metalloproteinase, MT4-MMP, MT4MMP, MMP17, MT4MMP

#### **Application Details**

WB 1:500-1:1000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200

#### **Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

# **Immunogen**

A synthesized peptide derived from human MMP17

#### **Purification**

Affinity-chromatography

Storage Store at -20°C for one year. For short term

storage and frequent use, store at 4°C for

up to one month. Avoid repeated

freeze-thaw cycles.

#### Anti-MMP17 Rabbit Monoclonal Antibody - Protein Information

Name MMP17

Synonyms MT4MMP



#### **Function**

Endopeptidase that degrades various components of the extracellular matrix, such as fibrin. May be involved in the activation of membrane-bound precursors of growth factors or inflammatory mediators, such as tumor necrosis factor-alpha. May also be involved in tumoral process. Cleaves pro-TNF-alpha at the '74-Ala-|-Gln-75' site. Not obvious if able to proteolytically activate progelatinase A. Does not hydrolyze collagen types I, II, III, IV and V, gelatin, fibronectin, laminin, decorin nor alpha1-antitrypsin.

# **Cellular Location**

[Isoform Long]: Cell membrane; Lipid-anchor, GPI- anchor; Extracellular side. Secreted, extracellular space, extracellular matrix

#### **Tissue Location**

Expressed in brain, leukocytes, colon, ovary testis and breast cancer. Expressed also in many transformed and non-transformed cell types

### **Anti-MMP17 Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-MMP17 Rabbit Monoclonal Antibody - Images