

**Anti-DKC1 Rabbit Monoclonal Antibody**  
Catalog # ABO15266**Specification****Anti-DKC1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC
Primary Accession	<a href="#">O60832</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-DKC1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

**Anti-DKC1 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 1736

**Other Names**

H/ACA ribonucleoprotein complex subunit DKC1, 5.4.99.-, CBF5 homolog, Dyskerin, Nopp140-associated protein of 57 kDa, Nucleolar protein NAP57, Nucleolar protein family A member 4, snoRNP protein DKC1, DKC1 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=2890](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=2890))>HGNC:2890</a>), NOLA4

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human DKC1

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-DKC1 Rabbit Monoclonal Antibody - Protein Information**

**Name** DKC1 ([HGNC:2890](#))

## Synonyms NOLA4

### Function

[Isoform 1]: Catalytic subunit of H/ACA small nucleolar ribonucleoprotein (H/ACA snoRNP) complex, which catalyzes pseudouridylation of rRNA (PubMed: [25219674](http://www.uniprot.org/citations/25219674), PubMed: [32554502](http://www.uniprot.org/citations/32554502)). This involves the isomerization of uridine such that the ribose is subsequently attached to C5, instead of the normal N1 (PubMed: [25219674](http://www.uniprot.org/citations/25219674)). Each rRNA can contain up to 100 pseudouridine ('psi') residues, which may serve to stabilize the conformation of rRNAs. Required for ribosome biogenesis and telomere maintenance (PubMed: [19179534](http://www.uniprot.org/citations/19179534), PubMed: [25219674](http://www.uniprot.org/citations/25219674)). Also required for correct processing or intranuclear trafficking of TERC, the RNA component of the telomerase reverse transcriptase (TERT) holoenzyme (PubMed: [19179534](http://www.uniprot.org/citations/19179534)).

### Cellular Location

[Isoform 1]: Nucleus, nucleolus. Nucleus, Cajal body {ECO:0000250|UniProtKB:P40615}

### Tissue Location


Ubiquitously expressed.

## Anti-DKC1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-DKC1 Rabbit Monoclonal Antibody - Images



Western blot analysis of DKC1 expression in SW480 cell lysate.