

## Anti-SF3B1 Rabbit Monoclonal Antibody Catalog # ABO15284

### Specification

---

#### Anti-SF3B1 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	<a href="#">O75533</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

#### Description

Anti-SF3B1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

#### Anti-SF3B1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 23451

#### Other Names

Splicing factor 3B subunit 1, Pre-mRNA-splicing factor SF3b 155 kDa subunit, SF3b155 {ECO:0000303|Ref.39}, Spliceosome-associated protein 155, SAP 155, SF3B1 {ECO:0000303|PubMed:30567737, ECO:0000312|HGNC:HGNC:10768}

#### Application Details

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>FC 1:500

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human SF3B1

#### Purification

Affinity-chromatography

#### Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

#### Anti-SF3B1 Rabbit Monoclonal Antibody - Protein Information

**Name** SF3B1 {ECO:0000303|PubMed:30567737, ECO:0000312|HGNC:HGNC:10768}

#### Function

Component of the 17S U2 SnRNP complex of the spliceosome, a large ribonucleoprotein complex that removes introns from transcribed pre-mRNAs (PubMed:<a href="http://www.uniprot.org/citations/12234937" target="\_blank">12234937</a>, PubMed:<a href="http://www.uniprot.org/citations/27720643" target="\_blank">27720643</a>, PubMed:<a href="http://www.uniprot.org/citations/32494006" target="\_blank">32494006</a>, PubMed:<a href="http://www.uniprot.org/citations/34822310" target="\_blank">34822310</a>). The 17S U2 SnRNP complex (1) directly participates in early spliceosome assembly and (2) mediates recognition of the intron branch site during pre-mRNA splicing by promoting the selection of the pre-mRNA branch-site adenosine, the nucleophile for the first step of splicing (PubMed:<a href="http://www.uniprot.org/citations/32494006" target="\_blank">32494006</a>, PubMed:<a href="http://www.uniprot.org/citations/34822310" target="\_blank">34822310</a>). Within the 17S U2 SnRNP complex, SF3B1 is part of the SF3B subcomplex, which is required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence in pre-mRNA (PubMed:<a href="http://www.uniprot.org/citations/12234937" target="\_blank">12234937</a>). Sequence independent binding of SF3A and SF3B subcomplexes upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA (PubMed:<a href="http://www.uniprot.org/citations/12234937" target="\_blank">12234937</a>). May also be involved in the assembly of the 'E' complex (PubMed:<a href="http://www.uniprot.org/citations/10882114" target="\_blank">10882114</a>). Also acts as a component of the minor spliceosome, which is involved in the splicing of U12-type introns in pre-mRNAs (PubMed:<a href="http://www.uniprot.org/citations/15146077" target="\_blank">15146077</a>, PubMed:<a href="http://www.uniprot.org/citations/33509932" target="\_blank">33509932</a>). Together with other U2 snRNP complex components may also play a role in the selective processing of microRNAs (miRNAs) from the long primary miRNA transcript, pri-miR-17-92 (By similarity).

#### **Cellular Location**

Nucleus. Nucleus speckle. Note=During mitosis, transiently dispersed from the nuclear speckles to the cytoplasm


#### **Anti-SF3B1 Rabbit Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti-SF3B1 Rabbit Monoclonal Antibody - Images**





Western blot analysis of SF3B1 expression in Jurkat cell lysate.