

Anti-MEIS2 Rabbit Monoclonal Antibody Catalog # ABO15285

Specification

Anti-MEIS2 Rabbit Monoclonal Antibody - Product Information

Application	WB, FC
Primary Accession	O14770
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-MEIS2 Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-MEIS2 Rabbit Monoclonal Antibody - Additional Information

Gene ID 4212

Other Names

Homeobox protein Meis2, Meis1-related protein 1, MEIS2, MRG1

Application Details

WB 1:500-1:2000
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human MEIS2

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-MEIS2 Rabbit Monoclonal Antibody - Protein Information

Name MEIS2

Synonyms MRG1

Function

Involved in transcriptional regulation. Binds to HOX or PBX proteins to form dimers, or to a DNA-bound dimer of PBX and HOX proteins and thought to have a role in stabilization of the homeoprotein-DNA complex. Isoform 3 is required for the activity of a PDX1:PBX1b:MEIS2b complex in pancreatic acinar cells involved in the transcriptional activation of the ELA1 enhancer; the complex binds to the enhancer B element and cooperates with the transcription factor 1 complex (PTF1) bound to the enhancer A element; MEIS2 is not involved in complex DNA-binding. Probably in complex with PBX1, is involved in transcriptional regulation by KLF4. Isoform 3 and isoform 4 can bind to a EPHA8 promoter sequence containing the DNA motif 5'-CGGTCA-3'; in cooperation with a PBX protein (such as PBX2) is proposed to be involved in the transcriptional activation of EPHA8 in the developing midbrain. May be involved in regulation of myeloid differentiation. Can bind to the DNA sequence 5'-TGACAG-3' in the activator ACT sequence of the D(1A) dopamine receptor (DRD1) promoter and activate DRD1 transcription; isoform 5 cannot activate DRD1 transcription.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108}. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:P97367}

Tissue Location


Expressed in various tissues. Expressed at high level in the lymphoid organs of hematopoietic tissues. Also expressed in some regions of the brain, such as the putamen

Anti-MEIS2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-MEIS2 Rabbit Monoclonal Antibody - Images



Western blot analysis of MEIS2 expression in HepG2 cell lysate.