

**Anti-hnRNP Q Rabbit Monoclonal Antibody**  
Catalog # ABO15327**Specification****Anti-hnRNP Q Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC
Primary Accession	<a href="#">O60506</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-hnRNP Q Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

**Anti-hnRNP Q Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 10492

**Other Names**

Heterogeneous nuclear ribonucleoprotein Q, hnRNP Q, Glycine- and tyrosine-rich RNA-binding protein, GRY-RBP, NS1-associated protein 1, Synaptotagmin-binding, cytoplasmic RNA-interacting protein, SYNCRIP, HNRPQ, NSAP1

**Calculated MW**

65-82 kDa KDa

**Application Details**

WB 1:1000-1:5000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human hnRNP Q

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-hnRNP Q Rabbit Monoclonal Antibody - Protein Information**

**Name** SYNCRIP

**Synonyms** HNRPQ, NSAP1

### Function

Heterogenous nuclear ribonucleoprotein (hnRNP) implicated in mRNA processing mechanisms. Component of the CRD-mediated complex that promotes MYC mRNA stability. Isoform 1, isoform 2 and isoform 3 are associated in vitro with pre-mRNA, splicing intermediates and mature mRNA protein complexes. Isoform 1 binds to apoB mRNA AU-rich sequences. Isoform 1 is part of the APOB mRNA editosome complex and may modulate the postranscriptional C to U RNA-editing of the APOB mRNA through either by binding to A1CF (APOBEC1 complementation factor), to APOBEC1 or to RNA itself. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain. Interacts in vitro preferentially with poly(A) and poly(U) RNA sequences. Isoform 3 may be involved in cytoplasmic vesicle-based mRNA transport through interaction with synaptotagmins. Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes. Upon interferon-gamma activation assembles into the GAIT complex which binds to stem loop- containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation; seems not to be essential for GAIT complex function.

### Cellular Location

Cytoplasm. Microsome {ECO:0000250|UniProtKB:Q7TMK9} Endoplasmic reticulum. Nucleus {ECO:0000250|UniProtKB:Q7TMK9}. Note=The tyrosine phosphorylated form bound to RNA is found in microsomes (By similarity). Localized in cytoplasmic mRNP granules containing untranslated mRNAs (By similarity). {ECO:0000250|UniProtKB:O43390, ECO:0000250|UniProtKB:Q7TMK9} [Isoform 2]: Nucleus, nucleoplasm {ECO:0000250|UniProtKB:Q7TMK9}. Note=Expressed predominantly in the nucleoplasm. {ECO:0000250|UniProtKB:Q7TMK9}

### Tissue Location


Ubiquitously expressed. Detected in heart, brain, pancreas, placenta, spleen, lung, liver, skeletal muscle, kidney, thymus, prostate, uterus, small intestine, colon, peripheral blood and testis.


## Anti-hnRNP Q Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-hnRNP Q Rabbit Monoclonal Antibody - Images






All lanes use the Antibody at 1:3W dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:3W dilution for 1 hour at room temperature.



Western blot analysis of hnRNP Q expression in HeLa cell lysate.