

Anti-IMPDH2 Rabbit Monoclonal Antibody
Catalog # ABO15471**Specification****Anti-IMPDH2 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P12268
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-IMPDH2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-IMPDH2 Rabbit Monoclonal Antibody - Additional Information

Gene ID 3615

Other Names

Inosine-5'-monophosphate dehydrogenase 2, IMP dehydrogenase 2, IMPD 2, IMPDH 2, 1.1.1.205, Inosine-5'-monophosphate dehydrogenase type II, IMP dehydrogenase II, IMPDH-II, IMPDH2 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=6053 target="_blank">HGNC:6053), IMPD2

Calculated MW

56 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:50
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human IMPDH2

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-IMPDH2 Rabbit Monoclonal Antibody - Protein Information

Name IMPDH2 ([HGNC:6053](#))

Synonyms IMPD2

Function

Catalyzes the conversion of inosine 5'-phosphate (IMP) to xanthosine 5'-phosphate (XMP), the first committed and rate-limiting step in the de novo synthesis of guanine nucleotides, and therefore plays an important role in the regulation of cell growth (PubMed:[7763314](http://www.uniprot.org/citations/7763314), PubMed:[7903306](http://www.uniprot.org/citations/7903306)). Could also have a single-stranded nucleic acid-binding activity and could play a role in RNA and/or DNA metabolism (PubMed:[14766016](http://www.uniprot.org/citations/14766016)). It may also have a role in the development of malignancy and the growth progression of some tumors.

Cellular Location

Cytoplasm. Nucleus. Cytoplasm, cytosol. Note=Can form fiber-like subcellular structures termed 'cytoophidia' in response to intracellular guanine- nucleotide depletion.

Tissue Location


IMPDH1 is the main species in normal leukocytes and IMPDH2 predominates over IMPDH1 in the tumor

Anti-IMPDH2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IMPDH2 Rabbit Monoclonal Antibody - Images



Western blot analysis of IMPDH2 expression in HeLa cell lysate.