

## Anti-LIMK1 Rabbit Monoclonal Antibody Catalog # ABO15487

### Specification

---

#### Anti-LIMK1 Rabbit Monoclonal Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P53667</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

#### Description

Anti-LIMK1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human Mouse Rat.

#### Anti-LIMK1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 3984

#### Other Names

LIM domain kinase 1, LIMK-1, 2.7.11.1, LIMK1, LIMK

#### Application Details

WB 1:500-1:2000

#### Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

#### Immunogen

A synthesized peptide derived from human LIM Kinase 1

#### Purification

Affinity-chromatography

Storage

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

#### Anti-LIMK1 Rabbit Monoclonal Antibody - Protein Information

Name LIMK1

Synonyms LIMK

Function

Serine/threonine-protein kinase that plays an essential role in the regulation of actin filament dynamics. Acts downstream of several Rho family GTPase signal transduction pathways (PubMed:<a href="http://www.uniprot.org/citations/10436159" target="\_blank">10436159</a>, PubMed:<a href="http://www.uniprot.org/citations/11832213" target="\_blank">11832213</a>, PubMed:<a href="http://www.uniprot.org/citations/12807904" target="\_blank">12807904</a>, PubMed:<a href="http://www.uniprot.org/citations/15660133" target="\_blank">15660133</a>, PubMed:<a href="http://www.uniprot.org/citations/16230460" target="\_blank">16230460</a>, PubMed:<a href="http://www.uniprot.org/citations/18028908" target="\_blank">18028908</a>, PubMed:<a href="http://www.uniprot.org/citations/22328514" target="\_blank">22328514</a>, PubMed:<a href="http://www.uniprot.org/citations/23633677" target="\_blank">23633677</a>). Activated by upstream kinases including ROCK1, PAK1 and PAK4, which phosphorylate LIMK1 on a threonine residue located in its activation loop (PubMed:<a href="http://www.uniprot.org/citations/10436159" target="\_blank">10436159</a>). LIMK1 subsequently phosphorylates and inactivates the actin binding/depolymerizing factors cofilin-1/CFL1, cofilin-2/CFL2 and destrin/DSTN, thereby preventing the cleavage of filamentous actin (F-actin), and stabilizing the actin cytoskeleton (PubMed:<a href="http://www.uniprot.org/citations/11832213" target="\_blank">11832213</a>, PubMed:<a href="http://www.uniprot.org/citations/15660133" target="\_blank">15660133</a>, PubMed:<a href="http://www.uniprot.org/citations/16230460" target="\_blank">16230460</a>, PubMed:<a href="http://www.uniprot.org/citations/23633677" target="\_blank">23633677</a>). In this way LIMK1 regulates several actin-dependent biological processes including cell motility, cell cycle progression, and differentiation (PubMed:<a href="http://www.uniprot.org/citations/11832213" target="\_blank">11832213</a>, PubMed:<a href="http://www.uniprot.org/citations/15660133" target="\_blank">15660133</a>, PubMed:<a href="http://www.uniprot.org/citations/16230460" target="\_blank">16230460</a>, PubMed:<a href="http://www.uniprot.org/citations/23633677" target="\_blank">23633677</a>). Phosphorylates TPPP on serine residues, thereby promoting microtubule disassembly (PubMed:<a href="http://www.uniprot.org/citations/18028908" target="\_blank">18028908</a>). Stimulates axonal outgrowth and may be involved in brain development (PubMed:<a href="http://www.uniprot.org/citations/18028908" target="\_blank">18028908</a>).

#### Cellular Location

Cytoplasm. Nucleus. Cytoplasm, cytoskeleton. Cell projection, lamellipodium {ECO:0000250|UniProtKB:P53668} Note=Predominantly found in the cytoplasm. Localizes in the lamellipodium in a CDC42BPA, CDC42BPB and FAM89B/LRAP25-dependent manner. {ECO:0000250|UniProtKB:P53668}

#### Tissue Location


Highest expression in both adult and fetal nervous system. Detected ubiquitously throughout the different regions of adult brain, with highest levels in the cerebral cortex. Expressed to a lesser extent in heart and skeletal muscle

### Anti-LIMK1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-LIMK1 Rabbit Monoclonal Antibody - Images



Western blot analysis of LIM Kinase 1 expression in U-87MG cell lysate.