

**Anti-U2AF65 Rabbit Monoclonal Antibody**  
Catalog # ABO15545

**Specification**

---

**Anti-U2AF65 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC
Primary Accession	<a href="#">P26368</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-U2AF65 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

**Anti-U2AF65 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 11338

**Other Names**

Splicing factor U2AF 65 kDa subunit, U2 auxiliary factor 65 kDa subunit, hU2AF(65), hU2AF65, U2 snRNP auxiliary factor large subunit, U2AF2, U2AF65

**Calculated MW**

65 kDa KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human U2AF65

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-U2AF65 Rabbit Monoclonal Antibody - Protein Information**

**Name** U2AF2

## Synonyms U2AF65

### Function

Plays a role in pre-mRNA splicing and 3'-end processing (PubMed:<a href="http://www.uniprot.org/citations/17024186" target="\_blank">17024186</a>). By recruiting PRPF19 and the PRP19C/Prp19 complex/NTC/Nineteen complex to the RNA polymerase II C-terminal domain (CTD), and thereby pre-mRNA, may couple transcription to splicing (PubMed:<a href="http://www.uniprot.org/citations/21536736" target="\_blank">21536736</a>). Induces cardiac troponin-T (TNNT2) pre-mRNA exon inclusion in muscle. Regulates the TNNT2 exon 5 inclusion through competition with MBNL1. Binds preferentially to a single-stranded structure within the polypyrimidine tract of TNNT2 intron 4 during spliceosome assembly. Required for the export of mRNA out of the nucleus, even if the mRNA is encoded by an intron-less gene. Represses the splicing of MAPT/Tau exon 10. Positively regulates pre-mRNA 3'-end processing by recruiting the CFIm complex to cleavage and polyadenylation signals (PubMed:<a href="http://www.uniprot.org/citations/17024186" target="\_blank">17024186</a>).

### Cellular Location


Nucleus.

## Anti-U2AF65 Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## Anti-U2AF65 Rabbit Monoclonal Antibody - Images



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



Western blot analysis of U2AF65 expression in Jurkat cell lysate.