

Anti-IFNGR1 Rabbit Monoclonal Antibody
Catalog # ABO15571**Specification****Anti-IFNGR1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, FC
Primary Accession	P15260
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-IFNGR1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.

Anti-IFNGR1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 3459

Other Names

Interferon gamma receptor 1 {ECO:0000312|HGNC:HGNC:5439}, IFN-gamma receptor 1, IFN-gamma-R1, CDw119, Interferon gamma receptor alpha-chain, IFN-gamma-R-alpha, CD119, IFNGR1 ([HGNC:5439](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=5439))

Calculated MW

45-100 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
FC 1:500

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human IFNGR1

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-IFNGR1 Rabbit Monoclonal Antibody - Protein Information

Name IFNGR1 ([HGNC:5439](#))

Function

Receptor subunit for interferon gamma/INFG that plays crucial roles in antimicrobial, antiviral, and antitumor responses by activating effector immune cells and enhancing antigen presentation (PubMed: [20015550](http://www.uniprot.org/citations/20015550)). Associates with transmembrane accessory factor IFNGR2 to form a functional receptor (PubMed: [10986460](http://www.uniprot.org/citations/10986460), PubMed: [2971451](http://www.uniprot.org/citations/2971451), PubMed: [7615558](http://www.uniprot.org/citations/7615558), PubMed: [7617032](http://www.uniprot.org/citations/7617032), PubMed: [7673114](http://www.uniprot.org/citations/7673114)). Upon ligand binding, the intracellular domain of IFNGR1 opens out to allow association of downstream signaling components JAK1 and JAK2. In turn, activated JAK1 phosphorylates IFNGR1 to form a docking site for STAT1. Subsequent phosphorylation of STAT1 leads to dimerization, translocation to the nucleus, and stimulation of target gene transcription (PubMed: [28883123](http://www.uniprot.org/citations/28883123)). STAT3 can also be activated in a similar manner although activation seems weaker. IFNGR1 intracellular domain phosphorylation also provides a docking site for SOCS1 that regulates the JAK-STAT pathway by competing with STAT1 binding to IFNGR1 (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein


Anti-IFNGR1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IFNGR1 Rabbit Monoclonal Antibody - Images





Western blot analysis of IFNGR1 expression in MCF7 cell lysate.