

Anti-ICOS Rabbit Monoclonal Antibody Catalog # ABO15765

Specification

Anti-ICOS Rabbit Monoclonal Antibody - Product Information

Application	WB, IF, ICC, FC
Primary Accession	O9Y6W8
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-ICOS Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Anti-ICOS Rabbit Monoclonal Antibody - Additional Information

Gene ID 29851

Other Names

Inducible T-cell costimulator, Activation-inducible lymphocyte immunomediatory molecule, CD278, ICOS, AILIM

Calculated MW

22 kDa KDa

Application Details

WB 1:500-1:2000
ICC/IF 1:50-1:200
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human ICOS

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-ICOS Rabbit Monoclonal Antibody - Protein Information

Name ICOS

Synonyms AILIM

Function

Enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells. Essential both for efficient interaction between T and B-cells and for normal antibody responses to T-cell dependent antigens. Does not up-regulate the production of interleukin- 2, but superinduces the synthesis of interleukin-10. Prevents the apoptosis of pre-activated T-cells. Plays a critical role in CD40- mediated class switching of immunoglobulin isotypes (By similarity).

Cellular Location

[Isoform 1]: Cell membrane; Single- pass type I membrane protein

Tissue Location


Activated T-cells. Highly expressed on tonsillar T- cells, which are closely associated with B-cells in the apical light zone of germinal centers, the site of terminal B-cell maturation Expressed at lower levels in thymus, lung, lymph node and peripheral blood leukocytes. Expressed in the medulla of fetal and newborn thymus

Anti-ICOS Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-ICOS Rabbit Monoclonal Antibody - Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of ICOS expression in Jurkat cell lysate.