

**Anti-TIP49A Rabbit Monoclonal Antibody**  
**Catalog # ABO15853****Specification**

---

**Anti-TIP49A Rabbit Monoclonal Antibody - Product Information**

Application	WB, FC
Primary Accession	<a href="#">Q9Y265</a>
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-TIP49A Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

**Anti-TIP49A Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 8607

**Other Names**

RuvB-like 1, 3.6.4.12, 49 kDa TATA box-binding protein-interacting protein, 49 kDa TBP-interacting protein, 54 kDa erythrocyte cytosolic protein, ECP-54, INO80 complex subunit H, Nuclear matrix protein 238, NMP 238, Pontin 52, TIP49a, TIP60-associated protein 54-alpha, TAP54-alpha, RUVBL1 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=10474](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=10474))  
HGNC:10474

**Calculated MW**

50 kDa KDa

**Application Details**

WB 1:500-1:2000  
FC 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human TIP49A

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-TIP49A Rabbit Monoclonal Antibody - Protein Information**

**Name** RUVBL1 ([HGNC:10474](#))

### Function

Possesses single-stranded DNA-stimulated ATPase and ATP- dependent DNA helicase (3' to 5') activity; hexamerization is thought to be critical for ATP hydrolysis and adjacent subunits in the ring- like structure contribute to the ATPase activity (PubMed:<a href="http://www.uniprot.org/citations/17157868" target="\_blank">17157868</a>, PubMed:<a href="http://www.uniprot.org/citations/33205750" target="\_blank">33205750</a>). Component of the NuA4 histone acetyltransferase complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A (PubMed:<a href="http://www.uniprot.org/citations/14966270" target="\_blank">14966270</a>). This modification may both alter nucleosome-DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription (PubMed:<a href="http://www.uniprot.org/citations/14966270" target="\_blank">14966270</a>). This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair (PubMed:<a href="http://www.uniprot.org/citations/14966270" target="\_blank">14966270</a>). The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. NuA4 may also play a direct role in DNA repair when recruited to sites of DNA damage (PubMed:<a href="http://www.uniprot.org/citations/14966270" target="\_blank">14966270</a>). Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AZ1 from the nucleosome (PubMed:<a href="http://www.uniprot.org/citations/24463511" target="\_blank">24463511</a>). Proposed core component of the chromatin remodeling INO80 complex which exhibits DNA- and nucleosome-activated ATPase activity and catalyzes ATP-dependent nucleosome sliding (PubMed:<a href="http://www.uniprot.org/citations/16230350" target="\_blank">16230350</a>, PubMed:<a href="http://www.uniprot.org/citations/21303910" target="\_blank">21303910</a>). Plays an essential role in oncogenic transformation by MYC and also modulates transcriptional activation by the LEF1/TCF1-CTNNB1 complex (PubMed:<a href="http://www.uniprot.org/citations/10882073" target="\_blank">10882073</a>, PubMed:<a href="http://www.uniprot.org/citations/16014379" target="\_blank">16014379</a>). Essential for cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/14506706" target="\_blank">14506706</a>). May be able to bind plasminogen at cell surface and enhance plasminogen activation (PubMed:<a href="http://www.uniprot.org/citations/11027681" target="\_blank">11027681</a>).

### Cellular Location

Nucleus matrix. Nucleus, nucleoplasm. Cytoplasm. Membrane Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Dynein axonemal particle {ECO:0000250|UniProtKB:Q9DE26}. Note=Mainly localized in the nucleus, associated with nuclear matrix or in the nuclear cytosol, although it is also present in the cytoplasm and associated with the cell membranes. In prophase and prometaphase it is located at the centrosome and the branching microtubule spindles. After mitotic nuclear membrane disintegration it accumulates at the centrosome and sites of tubulin polymerization. As cells pass through metaphase and into telophase it is located close to the centrosome at the early phase of tubulin polymerization. In anaphase it accumulates at the zone of tubule interdigitation. In telophase it is found at polar tubule overlap, and it reappears at the site of chromosomal decondensation in the daughter cells

### Tissue Location


Ubiquitously expressed with high expression in heart, skeletal muscle and testis

### Anti-TIP49A Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-TIP49A Rabbit Monoclonal Antibody - Images



Western blot analysis of TIP49A expression in K562 cell lysate.