

Anti-STK39 Rabbit Monoclonal Antibody
Catalog # ABO15938

Specification

Anti-STK39 Rabbit Monoclonal Antibody - Product Information

Application	WB, IHC, FC
Primary Accession	Q9UEW8
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-STK39 Rabbit Monoclonal Antibody . Tested in WB, IHC, Flow Cytometry applications. This antibody reacts with Human.

Anti-STK39 Rabbit Monoclonal Antibody - Additional Information

Gene ID 27347

Other Names

STE20/SPS1-related proline-alanine-rich protein kinase, Ste-20-related kinase, 2.7.11.1, DCHT, Serine/threonine-protein kinase 39, STK39

Calculated MW

65-70 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200
FC 1:50

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human STK39

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-STK39 Rabbit Monoclonal Antibody - Protein Information

Name STK39

Function

Effector serine/threonine-protein kinase component of the WNK-SPAK/OSR1 kinase cascade, which is involved in various processes, such as ion transport, response to hypertonic stress and blood pressure (PubMed: [16669787](http://www.uniprot.org/citations/16669787)), PubMed: [18270262](http://www.uniprot.org/citations/18270262), PubMed: [21321328](http://www.uniprot.org/citations/21321328), PubMed: [34289367](http://www.uniprot.org/citations/34289367)). Specifically recognizes and binds proteins with a RFXV motif (PubMed: [16669787](http://www.uniprot.org/citations/16669787)), PubMed: [21321328](http://www.uniprot.org/citations/21321328)). Acts downstream of WNK kinases (WNK1, WNK2, WNK3 or WNK4): following activation by WNK kinases, catalyzes phosphorylation of ion cotransporters, such as SLC12A1/NKCC2, SLC12A2/NKCC1, SLC12A3/NCC, SLC12A5/KCC2 or SLC12A6/KCC3, regulating their activity (PubMed: [21321328](http://www.uniprot.org/citations/21321328)). Mediates regulatory volume increase in response to hyperosmotic stress by catalyzing phosphorylation of ion cotransporters SLC12A1/NKCC2, SLC12A2/NKCC1 and SLC12A6/KCC3 downstream of WNK1 and WNK3 kinases (PubMed: [12740379](http://www.uniprot.org/citations/12740379)), PubMed: [16669787](http://www.uniprot.org/citations/16669787), PubMed: [21321328](http://www.uniprot.org/citations/21321328)). Phosphorylation of Na-K-Cl cotransporters SLC12A2/NKCC1 and SLC12A2/NKCC1 promote their activation and ion influx; simultaneously, phosphorylation of K-Cl cotransporters SLC12A5/KCC2 and SLC12A6/KCC3 inhibit their activity, blocking ion efflux (PubMed: [16669787](http://www.uniprot.org/citations/16669787)), PubMed: [19665974](http://www.uniprot.org/citations/19665974), PubMed: [21321328](http://www.uniprot.org/citations/21321328)). Acts as a regulator of NaCl reabsorption in the distal nephron by mediating phosphorylation and activation of the thiazide-sensitive Na-Cl cotransporter SLC12A3/NCC in distal convoluted tubule cells of kidney downstream of WNK4 (PubMed: [18270262](http://www.uniprot.org/citations/18270262)). Mediates the inhibition of SLC4A4, SLC26A6 as well as CFTR activities (By similarity). Phosphorylates RELT (By similarity).

Cellular Location

Cytoplasm. Nucleus. Note=Nucleus when caspase-cleaved.

Tissue Location


Predominantly expressed in brain and pancreas followed by heart, lung, kidney, skeletal muscle, liver, placenta and testis.


Anti-STK39 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-STK39 Rabbit Monoclonal Antibody - Images







All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



Western blot analysis of STK39 expression in HepG2 cell lysate.