

**Anti-TRAF2 Rabbit Monoclonal Antibody**  
**Catalog # ABO16050****Specification**

---

**Anti-TRAF2 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	<a href="#">Q12933</a>
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-TRAF2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.

**Anti-TRAF2 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 7186

**Other Names**

TNF receptor-associated factor 2, 2.3.2.27, E3 ubiquitin-protein ligase TRAF2, RING-type E3 ubiquitin transferase TRAF2, Tumor necrosis factor type 2 receptor-associated protein 3, TRAF2, TRAP3

**Calculated MW**

56 kDa KDa

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>IP 1:50<br>FC 1:50

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human TRAF2

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-TRAF2 Rabbit Monoclonal Antibody - Protein Information**

**Name** TRAF2 {ECO:0000303|PubMed:28489822, ECO:0000312|HGNC:HGNC:12032}

### Function

E3 ubiquitin-protein ligase that regulates activation of NF- kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/10346818" target="\_blank">10346818</a>, PubMed:<a href="http://www.uniprot.org/citations/11784851" target="\_blank">11784851</a>, PubMed:<a href="http://www.uniprot.org/citations/12917689" target="\_blank">12917689</a>, PubMed:<a href="http://www.uniprot.org/citations/15383523" target="\_blank">15383523</a>, PubMed:<a href="http://www.uniprot.org/citations/18981220" target="\_blank">18981220</a>, PubMed:<a href="http://www.uniprot.org/citations/19150425" target="\_blank">19150425</a>, PubMed:<a href="http://www.uniprot.org/citations/19810754" target="\_blank">19810754</a>, PubMed:<a href="http://www.uniprot.org/citations/19918265" target="\_blank">19918265</a>, PubMed:<a href="http://www.uniprot.org/citations/19937093" target="\_blank">19937093</a>, PubMed:<a href="http://www.uniprot.org/citations/20047764" target="\_blank">20047764</a>, PubMed:<a href="http://www.uniprot.org/citations/20064526" target="\_blank">20064526</a>, PubMed:<a href="http://www.uniprot.org/citations/20385093" target="\_blank">20385093</a>, PubMed:<a href="http://www.uniprot.org/citations/20577214" target="\_blank">20577214</a>, PubMed:<a href="http://www.uniprot.org/citations/22212761" target="\_blank">22212761</a>). Catalyzes 'Lys-63'-linked ubiquitination of target proteins, such as BIRC3, IKBKE, MLST8, RIPK1 and TICAM1 (PubMed:<a href="http://www.uniprot.org/citations/23453969" target="\_blank">23453969</a>, PubMed:<a href="http://www.uniprot.org/citations/28489822" target="\_blank">28489822</a>). Is an essential constituent of several E3 ubiquitin- protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases (PubMed:<a href="http://www.uniprot.org/citations/15383523" target="\_blank">15383523</a>, PubMed:<a href="http://www.uniprot.org/citations/18981220" target="\_blank">18981220</a>). Regulates BIRC2 and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain (PubMed:<a href="http://www.uniprot.org/citations/11907583" target="\_blank">11907583</a>, PubMed:<a href="http://www.uniprot.org/citations/19506082" target="\_blank">19506082</a>). Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR (PubMed:<a href="http://www.uniprot.org/citations/15121867" target="\_blank">15121867</a>). In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE (PubMed:<a href="http://www.uniprot.org/citations/23453969" target="\_blank">23453969</a>). Acts as a regulator of mTORC1 and mTORC2 assembly by mediating 'Lys-63'-linked ubiquitination of MLST8, thereby inhibiting formation of the mTORC2 complex, while facilitating assembly of the mTORC1 complex (PubMed:<a href="http://www.uniprot.org/citations/28489822" target="\_blank">28489822</a>). Required for normal antibody isotype switching from IgM to IgG (By similarity).

### Cellular Location


Cytoplasm

### Anti-TRAF2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-TRAF2 Rabbit Monoclonal Antibody - Images



Western blot analysis of TRAF2 expression in Raji cell lysate.