

Anti-IKZF3 Rabbit Monoclonal Antibody

Catalog # ABO16227

Specification

Anti-IKZF3 Rabbit Monoclonal Antibody - Product Information

Application WB, IHC
Primary Accession O9UKT9
Host Rabbit
Isotype IgG

Reactivity Rat, Human, Mouse

Clonality Monoclonal Format Liquid

Description

Anti-IKZF3 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

Anti-IKZF3 Rabbit Monoclonal Antibody - Additional Information

Gene ID 22806

Other Names

Zinc finger protein Aiolos, Ikaros family zinc finger protein 3, IKZF3, ZNFN1A3

Calculated MW 65 kDa KDa

Application Details

WB 1:500-1:2000
IHC 1:50-1:200

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human IKZF3

Purification

Affinity-chromatography

Storage Store at -20°C for one year. For short term

storage and frequent use, store at 4°C for

up to one month. Avoid repeated

freeze-thaw cycles.

Anti-IKZF3 Rabbit Monoclonal Antibody - Protein Information

Name IKZF3



Synonyms ZNFN1A3

Function

Transcription factor that plays an important role in the regulation of lymphocyte differentiation. Plays an essential role in regulation of B-cell differentiation, proliferation and maturation to an effector state. Involved in regulating BCL2 expression and controlling apoptosis in T-cells in an IL2-dependent manner.

Cellular Location

Nucleus. Cytoplasm [Isoform 3]: Nucleus [Isoform 14]: Nucleus. Cytoplasm

Tissue Location


Expressed most strongly in peripheral blood leukocytes, the spleen, and the thymus.

Anti-IKZF3 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-IKZF3 Rabbit Monoclonal Antibody - Images



Western blot analysis of IKZF3 expression in Raji cell lysate.