

**Anti-PCBP2/hnRNP E2 Antibody Picoband™ (monoclonal, 4B9C7)**  
Catalog # ABO16244**Specification****Anti-PCBP2/hnRNP E2 Antibody Picoband™ (monoclonal, 4B9C7) - Product Information**

Application	WB, IF, ICC, FC
Primary Accession	<a href="#">Q15366</a>
Host	Mouse
Isotype	Mouse IgG2a
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Lyophilized

**Description**

Anti-PCBP2/hnRNP E2 Antibody Picoband™ (monoclonal, 4B9C7) . Tested in Flow Cytometry, IF, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.

**Reconstitution**

Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

**Anti-PCBP2/hnRNP E2 Antibody Picoband™ (monoclonal, 4B9C7) - Additional Information**

Gene ID 5094

**Other Names**

Poly(rC)-binding protein 2, Alpha-CP2, Heterogeneous nuclear ribonucleoprotein E2, hnRNP E2, PCBP2 {ECO:0000303|PubMed:7607214, ECO:0000312|HGNC:HGNC:8648}

**Calculated MW**

39 kDa KDa

**Application Details**

Western blot, 0.25-0.5 µg/ml, Human, Mouse, Rat  
Immunocytochemistry/Immunofluorescence, 5 µg/ml, Human  
Flow Cytometry, 1-3 µg/1x10<sup>6</sup> cells, Human

**Contents**

Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na<sub>2</sub>HPO<sub>4</sub>.

**Immunogen**

E.coli-derived human PCBP2/hnRNP E2 recombinant protein (Position: Q197-K276).

**Purification**

Immunogen affinity purified.

**Storage**

**At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.**

## Anti-PCBP2/hnRNP E2 Antibody Picoband™ (monoclonal, 4B9C7) - Protein Information

**Name** PCBP2 {ECO:0000303|PubMed:7607214, ECO:0000312|HGNC:HGNC:8648}

### Function

Single-stranded nucleic acid binding protein that binds preferentially to oligo dC (PubMed:<a href="http://www.uniprot.org/citations/12414943" target="\_blank">12414943</a>, PubMed:<a href="http://www.uniprot.org/citations/7607214" target="\_blank">7607214</a>). Major cellular poly(rC)-binding protein (PubMed:<a href="http://www.uniprot.org/citations/12414943" target="\_blank">12414943</a>). Binds also poly(rU) (PubMed:<a href="http://www.uniprot.org/citations/12414943" target="\_blank">12414943</a>). Acts as a negative regulator of antiviral signaling (PubMed:<a href="http://www.uniprot.org/citations/19881509" target="\_blank">19881509</a>, PubMed:<a href="http://www.uniprot.org/citations/35322803" target="\_blank">35322803</a>). Negatively regulates cellular antiviral responses mediated by MAVS signaling (PubMed:<a href="http://www.uniprot.org/citations/19881509" target="\_blank">19881509</a>). It acts as an adapter between MAVS and the E3 ubiquitin ligase ITCH, therefore triggering MAVS ubiquitination and degradation (PubMed:<a href="http://www.uniprot.org/citations/19881509" target="\_blank">19881509</a>). Negatively regulates the cGAS-STING pathway via interaction with CGAS, preventing the formation of liquid- like droplets in which CGAS is activated (PubMed:<a href="http://www.uniprot.org/citations/35322803" target="\_blank">35322803</a>). Together with PCBP1, required for erythropoiesis, possibly by regulating mRNA splicing (By similarity).

### Cellular Location

Nucleus. Cytoplasm. Note=Loosely bound in the nucleus (PubMed:7607214). May shuttle between the nucleus and the cytoplasm (PubMed:7607214).

### Tissue Location


Detected in all tissues examined.

## Anti-PCBP2/hnRNP E2 Antibody Picoband™ (monoclonal, 4B9C7) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-PCBP2/hnRNP E2 Antibody Picoband™ (monoclonal, 4B9C7) - Images






Figure 1. Western blot analysis of PCBP2/hnRNP E2 using anti-PCBP2/hnRNP E2 antibody (M02425).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

- Lane 1: human T-47D whole cell lysates,
- Lane 2: human HL-60 whole cell lysates,
- Lane 3: human A549 whole cell lysates,
- Lane 4: human HepG2 whole cell lysates,
- Lane 5: rat brain tissue lysates,
- Lane 6: mouse brain tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-PCBP2/hnRNP E2 antigen affinity purified monoclonal antibody (Catalog # M02425) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for PCBP2/hnRNP E2 at approximately 39 kDa. The expected band size for PCBP2/hnRNP E2 is at 39 kDa.




Figure 2. IF analysis of PCBP2/hnRNP E2 using anti-PCBP2/hnRNP E2 antibody (M02425).

PCBP2/hnRNP E2 was detected in an immunocytochemical section of Caco-2 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 µg/mL mouse anti-PCBP2/hnRNP E2 Antibody (M02425) overnight at 4°C. DyLight®594 Conjugated Goat Anti-Mouse IgG (BA1141) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 3. Flow Cytometry analysis of PC-3 cells using anti-PCBP2/hnRNP E2 antibody (M02425). Overlay histogram showing PC-3 cells stained with M02425 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-PCBP2/hnRNP E2 Antibody (M02425, 1  $\mu\text{g}/1 \times 10^6$  cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10  $\mu\text{g}/1 \times 10^6$  cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1  $\mu\text{g}/1 \times 10^6$ ) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

### Anti-PCBP2/hnRNP E2 Antibody Picoband™ (monoclonal, 4B9C7) - Background

Poly(rC)-binding protein 2 is a protein that in humans is encoded by the PCBP2 gene. The protein encoded by this gene appears to be multifunctional. Along with PCBP-1 and hnRNPk, it is one of the major cellular poly(rC)-binding proteins. The encoded protein contains three K-homologous (KH) domains which may be involved in RNA binding. Together with PCBP-1, this protein also functions as a translational coactivator of poliovirus RNA via a sequence-specific interaction with stem-loop IV of the IRES, promoting poliovirus RNA replication by binding to its 5'-terminal cloverleaf structure. It has also been implicated in translational control of the 15-lipoxygenase mRNA, human papillomavirus type 16 L2 mRNA, and hepatitis A virus RNA. The encoded protein is also suggested to play a part in formation of a sequence-specific alpha-globin mRNP complex which is associated with alpha-globin mRNA stability. This multiexon structural mRNA is thought to be retrotransposed to generate PCBP-1, an intronless gene with functions similar to that of PCBP2. This gene and PCBP-1 have paralogous genes (PCBP3 and PCBP4) which are thought to have arisen as a result of duplication events of entire genes. This gene also has two processed pseudogenes (PCBP2P1 and PCBP2P2). Multiple transcript variants encoding different isoforms have been found for this gene.