

Anti-Nucleolin Rabbit Monoclonal Antibody
Catalog # ABO16369**Specification**

Anti-Nucleolin Rabbit Monoclonal Antibody - Product Information

Application	WB
Primary Accession	P19338
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

Description

Anti-Nucleolin Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.

Anti-Nucleolin Rabbit Monoclonal Antibody - Additional Information

Gene ID 4691

Other Names

Nucleolin, Protein C23, NCL

Calculated MW

100-110 kDa KDa

Application Details

WB 1:500-1:2000

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Nucleolin

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Nucleolin Rabbit Monoclonal Antibody - Protein Information

Name NCL

Function

Nucleolin is the major nucleolar protein of growing eukaryotic cells. It is found associated with intranucleolar chromatin and pre-ribosomal particles. It induces chromatin decondensation by binding to histone H1. It is thought to play a role in pre-rRNA transcription and ribosome assembly. May play a role in the process of transcriptional elongation. Binds RNA oligonucleotides with 5'-UUAGGG- 3' repeats more tightly than the telomeric single-stranded DNA 5'- TTAGGG-3' repeats.

Cellular Location


Nucleus, nucleolus. Cytoplasm. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs

Anti-Nucleolin Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Nucleolin Rabbit Monoclonal Antibody - Images



Western blot analysis of Nucleolin expression in HeLa cell lysate.