

Anti-XIAP Antibody Picoband™ (monoclonal, 3G2G1)
Catalog # ABO16608

Specification

Anti-XIAP Antibody Picoband™ (monoclonal, 3G2G1) - Product Information

Application	WB
Primary Accession	P98170
Host	Mouse
Isotype	Mouse IgG1
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-XIAP Antibody Picoband™ (monoclonal, 3G2G1) . Tested in WB applications. This antibody reacts with Human, Mouse, Rat.

Reconstitution

Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

Anti-XIAP Antibody Picoband™ (monoclonal, 3G2G1) - Additional Information

Gene ID 331

Other Names

E3 ubiquitin-protein ligase XIAP, 2.3.2.27, Baculoviral IAP repeat-containing protein 4, IAP-like protein, ILP, hILP, Inhibitor of apoptosis protein 3, IAP-3, hIAP-3, hIAP3, RING-type E3 ubiquitin transferase XIAP, X-linked inhibitor of apoptosis protein, X-linked IAP, XIAP {ECO:0000303|PubMed:12121969, ECO:0000312|HGNC:HGNC:592}

Calculated MW

54 kDa KDa

Application Details

Western blot, 0.25-0.5 µg/ml, Human, Mouse, Rat

Contents

Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na₂HPO₄.

Immunogen

E.coli-derived human XIAP recombinant protein (Position: A15-V244). Human XIAP shares 89.4% and 90.7% amino acid (aa) sequence identity with mouse and rat XIAP, respectively.

Purification

Immunogen affinity purified.

Storage

At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated

freezing and thawing.

Anti-XIAP Antibody Picoband™ (monoclonal, 3G2G1) - Protein Information

Name XIAP {ECO:0000303|PubMed:12121969, ECO:0000312|HGNC:HGNC:592}

Function

Multi-functional protein which regulates not only caspases and apoptosis, but also modulates inflammatory signaling and immunity, copper homeostasis, mitogenic kinase signaling, cell proliferation, as well as cell invasion and metastasis (PubMed: [11257230](http://www.uniprot.org/citations/11257230), PubMed: [11257231](http://www.uniprot.org/citations/11257231), PubMed: [11447297](http://www.uniprot.org/citations/11447297), PubMed: [12121969](http://www.uniprot.org/citations/12121969), PubMed: [12620238](http://www.uniprot.org/citations/12620238), PubMed: [17560374](http://www.uniprot.org/citations/17560374), PubMed: [17967870](http://www.uniprot.org/citations/17967870), PubMed: [19473982](http://www.uniprot.org/citations/19473982), PubMed: [20154138](http://www.uniprot.org/citations/20154138), PubMed: [22103349](http://www.uniprot.org/citations/22103349), PubMed: [9230442](http://www.uniprot.org/citations/9230442)). Acts as a direct caspase inhibitor (PubMed: [11257230](http://www.uniprot.org/citations/11257230), PubMed: [11257231](http://www.uniprot.org/citations/11257231), PubMed: [12620238](http://www.uniprot.org/citations/12620238), PubMed: [12620238](http://www.uniprot.org/citations/12620238)). Directly bind to the active site pocket of CASP3 and CASP7 and obstructs substrate entry (PubMed: [11257230](http://www.uniprot.org/citations/11257230), PubMed: [11257231](http://www.uniprot.org/citations/11257231), PubMed: [16352606](http://www.uniprot.org/citations/16352606), PubMed: [16916640](http://www.uniprot.org/citations/16916640), PubMed: [16916640](http://www.uniprot.org/citations/16916640)). Inactivates CASP9 by keeping it in a monomeric, inactive state (PubMed: [12620238](http://www.uniprot.org/citations/12620238)). Acts as an E3 ubiquitin-protein ligase regulating NF-kappa-B signaling and the target proteins for its E3 ubiquitin-protein ligase activity include: RIPK1, RIPK2, MAP3K2/MEKK2, DIABLO/SMAC, AIFM1, CCS, PTEN and BIRC5/survivin (PubMed: [17560374](http://www.uniprot.org/citations/17560374), PubMed: [17967870](http://www.uniprot.org/citations/17967870), PubMed: [19473982](http://www.uniprot.org/citations/19473982), PubMed: [20154138](http://www.uniprot.org/citations/20154138), PubMed: [22103349](http://www.uniprot.org/citations/22103349), PubMed: [22607974](http://www.uniprot.org/citations/22607974), PubMed: [29452636](http://www.uniprot.org/citations/29452636), PubMed: [30026309](http://www.uniprot.org/citations/30026309)). Acts as an important regulator of innate immunity by mediating 'Lys-63'-linked polyubiquitination of RIPK2 downstream of NOD1 and NOD2, thereby transforming RIPK2 into a scaffolding protein for downstream effectors, ultimately leading to activation of the NF-kappa-B and MAP kinases signaling (PubMed: [19667203](http://www.uniprot.org/citations/19667203), PubMed: [22607974](http://www.uniprot.org/citations/22607974), PubMed: [22607974](http://www.uniprot.org/citations/22607974), PubMed: [29452636](http://www.uniprot.org/citations/29452636), PubMed: [29452636](http://www.uniprot.org/citations/29452636), PubMed: [30026309](http://www.uniprot.org/citations/30026309), PubMed: [30026309](http://www.uniprot.org/citations/30026309)). 'Lys-63'-linked polyubiquitination of RIPK2 also promotes recruitment of the LUBAC complex to RIPK2 (PubMed: [22607974](http://www.uniprot.org/citations/22607974), PubMed: [22607974](http://www.uniprot.org/citations/22607974), PubMed: [29452636](http://www.uniprot.org/citations/29452636), PubMed: [29452636](http://www.uniprot.org/citations/29452636)). Regulates the BMP signaling pathway and the SMAD and MAP3K7/TAK1 dependent pathways leading to NF-kappa-B and JNK activation (PubMed: [17560374](http://www.uniprot.org/citations/17560374), PubMed: [17560374](http://www.uniprot.org/citations/17560374)). Ubiquitination of CCS leads to enhancement of its chaperone

activity toward its physiologic target, SOD1, rather than proteasomal degradation (PubMed:20154138). Ubiquitination of MAP3K2/MEKK2 and AIFM1 does not lead to proteasomal degradation (PubMed:17967870, PubMed:22103349). Plays a role in copper homeostasis by ubiquitinating COMMD1 and promoting its proteasomal degradation (PubMed:14685266). Can also function as E3 ubiquitin-protein ligase of the NEDD8 conjugation pathway, targeting effector caspases for neddylation and inactivation (PubMed:21145488). Ubiquitinates and therefore mediates the proteasomal degradation of BCL2 in response to apoptosis (PubMed:29020630). Protects cells from spontaneous formation of the ripoptosome, a large multi-protein complex that has the capability to kill cancer cells in a caspase-dependent and caspase-independent manner (PubMed:22095281). Suppresses ripoptosome formation by ubiquitinating RIPK1 and CASP8 (PubMed:22095281). Acts as a positive regulator of Wnt signaling and ubiquitinates TLE1, TLE2, TLE3, TLE4 and AES (PubMed:22304967). Ubiquitination of TLE3 results in inhibition of its interaction with TCF7L2/TCF4 thereby allowing efficient recruitment and binding of the transcriptional coactivator beta-catenin to TCF7L2/TCF4 that is required to initiate a Wnt-specific transcriptional program (PubMed:22304967).

Cellular Location

Cytoplasm. Nucleus. Note=TLE3 promotes its nuclear localization.

Tissue Location


Expressed in colonic crypts (at protein level) (PubMed:30389919). Ubiquitous, except peripheral blood leukocytes (PubMed:8654366).

Anti-XIAP Antibody Picoband™ (monoclonal, 3G2G1) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-XIAP Antibody Picoband™ (monoclonal, 3G2G1) - Images






Figure 1. Western blot analysis of XIAP using anti-XIAP antibody (M00482-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

- Lane 1: human HeLa whole cell lysates,
- Lane 2: human MCF-7 whole cell lysates,
- Lane 3: human PANC-1 whole cell lysates,
- Lane 4: human U87 whole cell lysates,
- Lane 5: rat heart tissue lysates,
- Lane 6: rat testis tissue lysates,
- Lane 7: mouse heart tissue lysates,
- Lane 8: mouse testis tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-XIAP antigen affinity purified monoclonal antibody (Catalog # M00482-1) at 0.5 μ g/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for XIAP at approximately 54 kDa. The expected band size for XIAP is at 57 kDa.

Anti-XIAP Antibody Picoband™ (monoclonal, 3G2G1) - Background

XIAP, also known as IAP3 or BIRC4, is a protein that stops apoptotic cell death. It is mapped to chromosome Xq25. This gene encodes a protein that belongs to a family of apoptotic suppressor proteins. Members of this family share a conserved motif termed, baculovirus IAP repeat, which is necessary for their anti-apoptotic function. This protein functions through binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2 and inhibits apoptosis induced by menadione, a potent inducer of free radicals, and interleukin 1-beta converting enzyme. This protein also inhibits at least two members of the caspase family of cell-death proteases, caspase-3 and caspase-7. Mutations in this gene are the cause of X-linked lymphoproliferative syndrome. Alternate splicing results in multiple transcript variants.