

Anti-PARK7/DJ1 Antibody Picoband™ (monoclonal, 4B10)
Catalog # ABO16610

Specification

Anti-PARK7/DJ1 Antibody Picoband™ (monoclonal, 4B10) - Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	Q99497
Host	Mouse
Isotype	IgG2b
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized

Description

Anti-PARK7/DJ1 Antibody Picoband™ (monoclonal, 4B10) . Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human.

Reconstitution

Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

Anti-PARK7/DJ1 Antibody Picoband™ (monoclonal, 4B10) - Additional Information

Gene ID 11315

Other Names

Parkinson disease protein 7, Maillard deglycase, Oncogene DJ1, Parkinsonism-associated deglycase {ECO:0000312|HGNC:HGNC:16369}, Protein DJ-1, DJ-1, Protein/nucleic acid deglycase DJ-1, 3.1.2.-, 3.5.1.-, 3.5.1.124, PARK7 (HGNC:16369)

Calculated MW

22 kDa KDa

Application Details

Western blot, 0.25-0.5 µg/ml, Human
 Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/ml, Human
 Immunocytochemistry/Immunofluorescence, 5 µg/ml, Human
 Flow Cytometry, 1-3 µg/1x10⁶ cells, Human

Contents

Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na₂HPO₄.

Immunogen

E.coli-derived human PARK7 recombinant protein (Position: A2-D189). Human PARK7 shares 91% amino acid (aa) sequence identity with both mouse and rat PARK7.

Purification

Immunogen affinity purified.

Storage

At -20°C for one year from date of receipt.

After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.

Anti-PARK7/DJ1 Antibody Picoband™ (monoclonal, 4B10) - Protein Information

Name PARK7 ([HGNC:16369](#))

Function

Multifunctional protein with controversial molecular function which plays an important role in cell protection against oxidative stress and cell death acting as oxidative stress sensor and redox-sensitive chaperone and protease (PubMed: [12796482](http://www.uniprot.org/citations/12796482) target="_blank">12796482, PubMed: [17015834](http://www.uniprot.org/citations/17015834) target="_blank">17015834, PubMed: [18711745](http://www.uniprot.org/citations/18711745) target="_blank">18711745, PubMed: [19229105](http://www.uniprot.org/citations/19229105) target="_blank">19229105, PubMed: [20304780](http://www.uniprot.org/citations/20304780) target="_blank">20304780, PubMed: [25416785](http://www.uniprot.org/citations/25416785) target="_blank">25416785, PubMed: [26995087](http://www.uniprot.org/citations/26995087) target="_blank">26995087, PubMed: [28993701](http://www.uniprot.org/citations/28993701) target="_blank">28993701). It is involved in neuroprotective mechanisms like the stabilization of NFE2L2 and PINK1 proteins, male fertility as a positive regulator of androgen signaling pathway as well as cell growth and transformation through, for instance, the modulation of NF-kappa-B signaling pathway (PubMed: [12612053](http://www.uniprot.org/citations/12612053) target="_blank">12612053, PubMed: [14749723](http://www.uniprot.org/citations/14749723) target="_blank">14749723, PubMed: [15502874](http://www.uniprot.org/citations/15502874) target="_blank">15502874, PubMed: [17015834](http://www.uniprot.org/citations/17015834) target="_blank">17015834, PubMed: [18711745](http://www.uniprot.org/citations/18711745) target="_blank">18711745, PubMed: [21097510](http://www.uniprot.org/citations/21097510) target="_blank">21097510). Has been described as a protein and nucleotide deglycase that catalyzes the deglycation of the Maillard adducts formed between amino groups of proteins or nucleotides and reactive carbonyl groups of glyoxals (PubMed: [25416785](http://www.uniprot.org/citations/25416785) target="_blank">25416785, PubMed: [28596309](http://www.uniprot.org/citations/28596309) target="_blank">28596309). But this function is rebutted by other works (PubMed: [27903648](http://www.uniprot.org/citations/27903648) target="_blank">27903648, PubMed: [31653696](http://www.uniprot.org/citations/31653696) target="_blank">31653696). As a protein deglycase, repairs methylglyoxal- and glyoxal-glycated proteins, and releases repaired proteins and lactate or glycolate, respectively. Deglycates cysteine, arginine and lysine residues in proteins, and thus reactivates these proteins by reversing glycation by glyoxals. Acts on early glycation intermediates (hemithioacetals and aminocarbinals), preventing the formation of advanced glycation endproducts (AGE) that cause irreversible damage (PubMed: [25416785](http://www.uniprot.org/citations/25416785) target="_blank">25416785, PubMed: [26995087](http://www.uniprot.org/citations/26995087) target="_blank">26995087, PubMed: [28013050](http://www.uniprot.org/citations/28013050) target="_blank">28013050). Also functions as a nucleotide deglycase able to repair glycated guanine in the free nucleotide pool (GTP, GDP, GMP, dGTP) and in DNA and RNA. Is thus involved in a major nucleotide repair system named guanine glycation repair (GG repair), dedicated to reversing methylglyoxal and glyoxal damage via nucleotide sanitization and direct nucleic acid repair (PubMed: [28596309](http://www.uniprot.org/citations/28596309) target="_blank">28596309). Protects histones from adduction by methylglyoxal, controls the levels of methylglyoxal- derived argininine modifications on chromatin (PubMed: [30150385](http://www.uniprot.org/citations/30150385) target="_blank">30150385). Able to remove the glycations and restore histone 3, histone glycation disrupts both local and global chromatin architecture by altering histone-DNA interactions as well as histone acetylation and ubiquitination levels (PubMed: [30150385](http://www.uniprot.org/citations/30150385) target="_blank">30150385, PubMed: [30894531](http://www.uniprot.org/citations/30894531) target="_blank">30894531).

target="_blank">30894531). Displays a very low glyoxalase activity that may reflect its deglycase activity (PubMed:22523093, PubMed:28993701, PubMed:31653696). Eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death (PubMed:16390825). Required for correct mitochondrial morphology and function as well as for autophagy of dysfunctional mitochondria (PubMed:16632486, PubMed:19229105). Plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking (PubMed:18711745). Regulates astrocyte inflammatory responses, may modulate lipid rafts-dependent endocytosis in astrocytes and neuronal cells (PubMed:23847046). In pancreatic islets, involved in the maintenance of mitochondrial reactive oxygen species (ROS) levels and glucose homeostasis in an age- and diet dependent manner. Protects pancreatic beta cells from cell death induced by inflammatory and cytotoxic setting (By similarity). Binds to a number of mRNAs containing multiple copies of GG or CC motifs and partially inhibits their translation but dissociates following oxidative stress (PubMed:18626009). Metal-binding protein able to bind copper as well as toxic mercury ions, enhances the cell protection mechanism against induced metal toxicity (PubMed:23792957). In macrophages, interacts with the NADPH oxidase subunit NCF1 to direct NADPH oxidase-dependent ROS production, and protects against sepsis (By similarity).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q99LX0}; Lipid-anchor {ECO:0000250|UniProtKB:Q99LX0}. Cytoplasm. Nucleus. Membrane raft {ECO:0000250|UniProtKB:O88767}. Mitochondrion. Endoplasmic reticulum. Note=Under normal conditions, located predominantly in the cytoplasm and, to a lesser extent, in the nucleus and mitochondrion. Translocates to the mitochondrion and subsequently to the nucleus in response to oxidative stress and exerts an increased cytoprotective effect against oxidative damage (PubMed:18711745). Detected in tau inclusions in brains from neurodegenerative disease patients (PubMed:14705119). Membrane raft localization in astrocytes and neuronal cells requires palmitoylation

Tissue Location

Highly expressed in pancreas, kidney, skeletal muscle, liver, testis and heart. Detected at slightly lower levels in placenta and brain (at protein level). Detected in astrocytes, Sertoli cells, spermatogonia, spermatids and spermatozoa. Expressed by pancreatic islets at higher levels than surrounding exocrine tissues (PubMed:22611253).

Anti-PARK7/DJ1 Antibody Picoband™ (monoclonal, 4B10) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PARK7/DJ1 Antibody Picoband™ (monoclonal, 4B10) - Images




Figure 1. Western blot analysis of PARK7/DJ1 using anti-PARK7/DJ1 antibody (M00757-4). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human U2OS whole cell lysates,
Lane 2: human HepG2 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-PARK7/DJ1 antigen affinity purified monoclonal antibody (Catalog # M00757-4) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for PARK7/DJ1 at approximately 22 kDa. The expected band size for PARK7/DJ1 is at 22 kDa.




Figure 2. IHC analysis of PARK7/DJ1 using anti-PARK7/DJ1 antibody (M00757-4). PARK7/DJ1 was detected in a paraffin-embedded section of human endometrial carcinoma tissue.

Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-PARK7/DJ1 Antibody (M00757-4) overnight at 4°C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit (Catalog # SV0001) with DAB as the chromogen.




Figure 3. IHC analysis of PARK7/DJ1 using anti-PARK7/DJ1 antibody (M00757-4). PARK7/DJ1 was detected in a paraffin-embedded section of human esophageal squamous carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-PARK7/DJ1 Antibody (M00757-4) overnight at 4°C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit (Catalog # SV0001) with DAB as the chromogen.




Figure 4. IHC analysis of PARK7/DJ1 using anti-PARK7/DJ1 antibody (M00757-4). PARK7/DJ1 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-PARK7/DJ1 Antibody (M00757-4) overnight at 4°C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit (Catalog # SV0001) with DAB as the chromogen.




Figure 5. IF analysis of PARK7/DJ1 using anti-PARK7/DJ1 antibody (M00757-4). PARK7/DJ1 was detected in an immunocytochemical section of HeLa cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 $\mu\text{g}/\text{mL}$ mouse anti-PARK7/DJ1 Antibody (M00757-4) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 6. Flow Cytometry analysis of HeLa cells using anti-PARK7/DJ1 antibody (M00757-4). Overlay histogram showing HeLa cells stained with M00757-4 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-PARK7/DJ1 Antibody (M00757-4, 1 $\mu\text{g}/1 \times 10^6$ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 $\mu\text{g}/1 \times 10^6$ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 $\mu\text{g}/1 \times 10^6$) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Anti-PARK7/DJ1 Antibody Picoband™ (monoclonal, 4B10) - Background

Parkinson disease (autosomal recessive, early onset) 7, also known as DJ1, is a protein which in humans is encoded by the PARK7 gene. PARK7 belongs to the peptidase C56 family of proteins. PARK7 is mapped to chromosome 1p36. It acts as a positive regulator of androgen receptor-dependent transcription. It is also involved in tumorigenesis and in maintaining mitochondrial homeostasis. This gene may also function as a redox-sensitive chaperone, as a sensor for oxidative stress, and it apparently protects neurons against oxidative stress and cell death. It has been found that PARK7 mutations that impair transcriptional coactivator function can render dopaminergic neurons vulnerable to apoptosis and may contribute to the pathogenesis of Parkinson disease.