

Anti-RIP2 Rabbit Monoclonal Antibody Catalog # ABO16631

Specification

Anti-RIP2 Rabbit Monoclonal Antibody - Product Information

Application	WB
Primary Accession	O43353
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

Description

Anti-RIP2 Rabbit Monoclonal Antibody . Tested in WB applications. This antibody reacts with Human, Mouse, Rat.

Anti-RIP2 Rabbit Monoclonal Antibody - Additional Information

Gene ID 8767

Other Names

Receptor-interacting serine/threonine-protein kinase 2, 2.7.11.1, CARD-containing interleukin-1 beta-converting enzyme-associated kinase, CARD-containing IL-1 beta ICE-kinase, RIP-like-interacting CLARP kinase, Receptor-interacting protein 2, RIP-2, Tyrosine-protein kinase RIPK2, 2.7.10.2, RIPK2 {ECO:0000303|PubMed:30026309, ECO:0000312|HGNC:HGNC:10020}

Application Details

WB 1:500-1:2000

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human RIP2

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-RIP2 Rabbit Monoclonal Antibody - Protein Information

Name RIPK2 {ECO:0000303|PubMed:30026309, ECO:0000312|HGNC:HGNC:10020}

Function

Serine/threonine/tyrosine-protein kinase that plays an essential role in modulation of innate and adaptive immune responses (PubMed: [14638696](http://www.uniprot.org/citations/14638696) target="_blank">14638696, PubMed: [17054981](http://www.uniprot.org/citations/17054981) target="_blank">17054981, PubMed: [21123652](http://www.uniprot.org/citations/21123652) target="_blank">21123652, PubMed: [28656966](http://www.uniprot.org/citations/28656966) target="_blank">28656966, PubMed: [9575181](http://www.uniprot.org/citations/9575181) target="_blank">9575181, PubMed: [9642260](http://www.uniprot.org/citations/9642260) target="_blank">9642260). Acts as a key effector of NOD1 and NOD2 signaling pathways: upon activation by bacterial peptidoglycans, NOD1 and NOD2 oligomerize and recruit RIPK2 via CARD-CARD domains, leading to the formation of RIPK2 filaments (PubMed: [17054981](http://www.uniprot.org/citations/17054981) target="_blank">17054981, PubMed: [17562858](http://www.uniprot.org/citations/17562858) target="_blank">17562858, PubMed: [21123652](http://www.uniprot.org/citations/21123652) target="_blank">21123652, PubMed: [22607974](http://www.uniprot.org/citations/22607974) target="_blank">22607974, PubMed: [28656966](http://www.uniprot.org/citations/28656966) target="_blank">28656966, PubMed: [29452636](http://www.uniprot.org/citations/29452636) target="_blank">29452636, PubMed: [30026309](http://www.uniprot.org/citations/30026309) target="_blank">30026309). Once recruited, RIPK2 autophosphorylates and undergoes 'Lys-63'-linked polyubiquitination by E3 ubiquitin ligases XIAP, BIRC2 and BIRC3, as well as 'Met-1'-linked (linear) polyubiquitination by the LUBAC complex, becoming a scaffolding protein for downstream effectors (PubMed: [22607974](http://www.uniprot.org/citations/22607974) target="_blank">22607974, PubMed: [28545134](http://www.uniprot.org/citations/28545134) target="_blank">28545134, PubMed: [29452636](http://www.uniprot.org/citations/29452636) target="_blank">29452636, PubMed: [30026309](http://www.uniprot.org/citations/30026309) target="_blank">30026309, PubMed: [30279485](http://www.uniprot.org/citations/30279485) target="_blank">30279485, PubMed: [30478312](http://www.uniprot.org/citations/30478312) target="_blank">30478312). 'Met-1'-linked polyubiquitin chains attached to RIPK2 recruit IKBKG/NEMO, which undergoes 'Lys-63'-linked polyubiquitination in a RIPK2-dependent process (PubMed: [17562858](http://www.uniprot.org/citations/17562858) target="_blank">17562858, PubMed: [22607974](http://www.uniprot.org/citations/22607974) target="_blank">22607974, PubMed: [29452636](http://www.uniprot.org/citations/29452636) target="_blank">29452636, PubMed: [30026309](http://www.uniprot.org/citations/30026309) target="_blank">30026309). 'Lys-63'-linked polyubiquitin chains attached to RIPK2 serve as docking sites for TAB2 and TAB3 and mediate the recruitment of MAP3K7/TAK1 to IKBKG/NEMO, inducing subsequent activation of IKBKB/IKK (PubMed: [18079694](http://www.uniprot.org/citations/18079694) target="_blank">18079694). In turn, NF-kappa-B is released from NF-kappa-B inhibitors and translocates into the nucleus where it activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis (PubMed: [18079694](http://www.uniprot.org/citations/18079694) target="_blank">18079694). The protein kinase activity is dispensable for the NOD1 and NOD2 signaling pathways (PubMed: [29452636](http://www.uniprot.org/citations/29452636) target="_blank">29452636, PubMed: [30026309](http://www.uniprot.org/citations/30026309) target="_blank">30026309). Contributes to the tyrosine phosphorylation of the guanine exchange factor ARHGEF2 through Src tyrosine kinase leading to NF-kappa-B activation by NOD2 (PubMed: [21887730](http://www.uniprot.org/citations/21887730) target="_blank">21887730). Also involved in adaptive immunity: plays a role during engagement of the T-cell receptor (TCR) in promoting BCL10 phosphorylation and subsequent NF-kappa-B activation (PubMed: [14638696](http://www.uniprot.org/citations/14638696) target="_blank">14638696). Plays a role in the inactivation of RHOA in response to NGFR signaling (PubMed: [26646181](http://www.uniprot.org/citations/26646181) target="_blank">26646181).

Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein. Endoplasmic reticulum. Note=Recruited to the cell membrane by NOD2 following stimulation by bacterial peptidoglycans

Tissue Location


Detected in heart, brain, placenta, lung, peripheral blood leukocytes, spleen, kidney, testis, prostate, pancreas and lymph node.

Anti-RIP2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-RIP2 Rabbit Monoclonal Antibody - Images



Western blot analysis of RIP2 in K562 cell lysate.