

**Anti-Caspase-6 Rabbit Monoclonal Antibody**  
Catalog # ABO16722**Specification****Anti-Caspase-6 Rabbit Monoclonal Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P55212</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Caspase-6 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

**Anti-Caspase-6 Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 839

**Other Names**

Caspase-6, CASP-6, CSP-6, 3.4.22.59, Apoptotic protease Mch-2, Caspase-6 subunit p18, Caspase-6 subunit p20, Caspase-6 subunit p11, Caspase-6 subunit p10, CASP6 ([HGNC:1507](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=1507))

**Application Details**

WB 1:500-1:2000<br>IHC 1:50-1:200

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Caspase-6

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-Caspase-6 Rabbit Monoclonal Antibody - Protein Information**

**Name** CASP6 ([HGNC:1507](#))

## Function

Cysteine protease that plays essential roles in programmed cell death, axonal degeneration, development and innate immunity (PubMed:<a href="http://www.uniprot.org/citations/19133298" target="\_blank">19133298</a>, PubMed:<a href="http://www.uniprot.org/citations/22858542" target="\_blank">22858542</a>, PubMed:<a href="http://www.uniprot.org/citations/27032039" target="\_blank">27032039</a>, PubMed:<a href="http://www.uniprot.org/citations/28864531" target="\_blank">28864531</a>, PubMed:<a href="http://www.uniprot.org/citations/30420425" target="\_blank">30420425</a>, PubMed:<a href="http://www.uniprot.org/citations/32298652" target="\_blank">32298652</a>, PubMed:<a href="http://www.uniprot.org/citations/8663580" target="\_blank">8663580</a>). Acts as a non- canonical executioner caspase during apoptosis: localizes in the nucleus and cleaves the nuclear structural protein NUMA1 and lamin A/LMNA thereby inducing nuclear shrinkage and fragmentation (PubMed:<a href="http://www.uniprot.org/citations/11953316" target="\_blank">11953316</a>, PubMed:<a href="http://www.uniprot.org/citations/17401638" target="\_blank">17401638</a>, PubMed:<a href="http://www.uniprot.org/citations/8663580" target="\_blank">8663580</a>, PubMed:<a href="http://www.uniprot.org/citations/9463409" target="\_blank">9463409</a>). Lamin-A/LMNA cleavage is required for chromatin condensation and nuclear disassembly during apoptotic execution (PubMed:<a href="http://www.uniprot.org/citations/11953316" target="\_blank">11953316</a>). Acts as a regulator of liver damage by promoting hepatocyte apoptosis: in absence of phosphorylation by AMP-activated protein kinase (AMPK), catalyzes cleavage of BID, leading to cytochrome c release, thereby participating in nonalcoholic steatohepatitis (PubMed:<a href="http://www.uniprot.org/citations/32029622" target="\_blank">32029622</a>). Cleaves PARK7/DJ-1 in cells undergoing apoptosis (By similarity). Involved in intrinsic apoptosis by mediating cleavage of RIPK1 (PubMed:<a href="http://www.uniprot.org/citations/22858542" target="\_blank">22858542</a>). Furthermore, cleaves many transcription factors such as NF-kappa-B and cAMP response element-binding protein/CREBBP (PubMed:<a href="http://www.uniprot.org/citations/10559921" target="\_blank">10559921</a>, PubMed:<a href="http://www.uniprot.org/citations/14657026" target="\_blank">14657026</a>). Cleaves phospholipid scramblase proteins XKR4 and XKR9 (By similarity). In addition to apoptosis, involved in different forms of programmed cell death (PubMed:<a href="http://www.uniprot.org/citations/32298652" target="\_blank">32298652</a>). Plays an essential role in defense against viruses by acting as a central mediator of the ZBP1-mediated pyroptosis, apoptosis, and necroptosis (PANoptosis), independently of its cysteine protease activity (PubMed:<a href="http://www.uniprot.org/citations/32298652" target="\_blank">32298652</a>). PANoptosis is a unique inflammatory programmed cell death, which provides a molecular scaffold that allows the interactions and activation of machinery required for inflammasome/pyroptosis, apoptosis and necroptosis (PubMed:<a href="http://www.uniprot.org/citations/32298652" target="\_blank">32298652</a>). Mechanistically, interacts with RIPK3 and enhances the interaction between RIPK3 and ZBP1, leading to ZBP1-mediated inflammasome activation and cell death (PubMed:<a href="http://www.uniprot.org/citations/32298652" target="\_blank">32298652</a>). Plays an essential role in axon degeneration during axon pruning which is the remodeling of axons during neurogenesis but not apoptosis (By similarity). Regulates B-cell programs both during early development and after antigen stimulation (By similarity).

## Cellular Location

Cytoplasm. Nucleus


## Anti-Caspase-6 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-Caspase-6 Rabbit Monoclonal Antibody - Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of Caspase-6 expression in (1) Mouse spleen lysate; (2) Rat kidney cell lysate.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.