

Anti-Phospho-eIF4EBP1 (T70) Rabbit Monoclonal Antibody Catalog # ABO16777

Specification

Anti-Phospho-eIF4EBP1 (T70) Rabbit Monoclonal Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | Q13541 |
| Host | Rabbit |
| Isotype | Rabbit IgG |
| Reactivity | Human |
| Clonality | Monoclonal |
| Format | Liquid |

Description

Anti-Phospho-eIF4EBP1 (T70) Rabbit Monoclonal Antibody . Tested in WB applications. This antibody reacts with Human.

Anti-Phospho-eIF4EBP1 (T70) Rabbit Monoclonal Antibody - Additional Information

Gene ID 1978

Other Names

Eukaryotic translation initiation factor 4E-binding protein 1, 4E-BP1, eIF4E-binding protein 1, Phosphorylated heat- and acid-stable protein regulated by insulin 1, PHAS-I, EIF4EBP1

Application Details

WB 1:500-1:2000

Contents

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

Immunogen

A synthesized peptide derived from human Phospho-eIF4EBP1 (T70)

Purification

Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Anti-Phospho-eIF4EBP1 (T70) Rabbit Monoclonal Antibody - Protein Information

Name EIF4EBP1

Function

Repressor of translation initiation that regulates EIF4E activity by preventing its assembly into the

EIF4F complex: hypophosphorylated form competes with EIF4G1/EIF4G3 and strongly binds to EIF4E, leading to repress translation. In contrast, hyperphosphorylated form dissociates from EIF4E, allowing interaction between EIF4G1/EIF4G3 and EIF4E, leading to initiation of translation. Mediates the regulation of protein translation by hormones, growth factors and other stimuli that signal through the MAP kinase and mTORC1 pathways.

Cellular Location


Cytoplasm. Nucleus. Note=Localization to the nucleus is unaffected by phosphorylation status.
{ECO:0000250|UniProtKB:Q60876}

Anti-Phospho-eIF4EBP1 (T70) Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Phospho-eIF4EBP1 (T70) Rabbit Monoclonal Antibody - Images



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.