

**Anti-Phospho-Caveolin-1 (Y14) Rabbit Monoclonal Antibody**  
Catalog # ABO16778**Specification**

---

**Anti-Phospho-Caveolin-1 (Y14) Rabbit Monoclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q03135</a>
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid

**Description**

Anti-Phospho-Caveolin-1 (Y14) Rabbit Monoclonal Antibody . Tested in WB applications. This antibody reacts with Human.

**Anti-Phospho-Caveolin-1 (Y14) Rabbit Monoclonal Antibody - Additional Information**

**Gene ID** 857

**Other Names**

Caveolin-1, CAV1, CAV

**Application Details**

WB 1:500-1:2000

**Contents**

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.

**Immunogen**

A synthesized peptide derived from human Phospho-Caveolin-1 (Y14)

**Purification**

Affinity-chromatography

**Storage**

**Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.**

**Anti-Phospho-Caveolin-1 (Y14) Rabbit Monoclonal Antibody - Protein Information**

**Name** CAV1

**Synonyms** CAV

**Function**

May act as a scaffolding protein within caveolar membranes (PubMed:<a href="http://www.uniprot.org/citations/11751885" target="\_blank">11751885</a>). Forms a stable heterooligomeric complex with CAV2 that targets to lipid rafts and drives caveolae formation. Mediates the recruitment of CAVIN proteins (CAVIN1/2/3/4) to the caveolae (PubMed:<a href="http://www.uniprot.org/citations/19262564" target="\_blank">19262564</a>). Interacts directly with G-protein alpha subunits and can functionally regulate their activity (By similarity). Involved in the costimulatory signal essential for T-cell receptor (TCR)-mediated T-cell activation. Its binding to DPP4 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner (PubMed:<a href="http://www.uniprot.org/citations/17287217" target="\_blank">17287217</a>). Recruits CTNNB1 to caveolar membranes and may regulate CTNNB1-mediated signaling through the Wnt pathway (By similarity). Negatively regulates TGF $\beta$ 1-mediated activation of SMAD2/3 by mediating the internalization of TGF $\beta$ 1 from membrane rafts leading to its subsequent degradation (PubMed:<a href="http://www.uniprot.org/citations/25893292" target="\_blank">25893292</a>). Binds 20(S)-hydroxycholesterol (20(S)-OHC) (By similarity).

#### Cellular Location

Golgi apparatus membrane; Peripheral membrane protein. Cell membrane; Peripheral membrane protein. Membrane, caveola; Peripheral membrane protein. Membrane raft. Golgi apparatus, trans-Golgi network {ECO:0000250|UniProtKB:P33724} Note=Colocalized with DPP4 in membrane rafts. Potential hairpin-like structure in the membrane. Membrane protein of caveolae

#### Tissue Location

Skeletal muscle, liver, stomach, lung, kidney and heart (at protein level). Expressed in the brain

### Anti-Phospho-Caveolin-1 (Y14) Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-Phospho-Caveolin-1 (Y14) Rabbit Monoclonal Antibody - Images