

**FGF-10 Antibody**  
**Rabbit Polyclonal Antibody**  
**Catalog # ABV10801**

**Specification**

---

**FGF-10 Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O15520</a>
Other Accession	<a href="#">AAM46926</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>23436</b>

**FGF-10 Antibody - Additional Information**

**Gene ID** 2255

**Application & Usage**

**Western blot analysis (0.5-4 µg/ml).  
However, the optimal conditions should be  
determined individually. Other applications  
have not been tested. Recombinant human  
FGF-10 can be used as a positive control.**

**Other Names**

FGF10, FGF-10, FGF 10, Fibroblast growth factor 10; FGF-10; FGF10, FGF 10

**Target/Specificity**

FGF-10

**Antibody Form**

Liquid

**Appearance**

Colorless liquid

**Formulation**

100 µg (0.5 mg/ml) affinity purified rabbit anti-human FGF-10 polyclonal antibody in phosphate buffered saline (PBS), pH 7.2, containing 30% glycerol, 0.5% BSA, and 0.01% thimerosal.

**Handling**

The antibody solution should be gently mixed before use.

**Reconstitution & Storage**

-20 °C

**Background Descriptions**

**Precautions**

FGF-10 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### **FGF-10 Antibody - Protein Information**

**Name** FGF10

**Function**

Plays an important role in the regulation of embryonic development, cell proliferation and cell differentiation. Required for normal branching morphogenesis. May play a role in wound healing.

**Cellular Location**

Secreted.

### **FGF-10 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **FGF-10 Antibody - Images**

### **FGF-10 Antibody - Background**

Fibroblast Growth Factor-10 (also called KGF-2) is a heparin binding growth factor that stimulates the proliferation and activation of cells that express FGF receptors. FGF-10 is mostly related to FGF-7/KGF and is expressed during development and preferentially in adult lungs.