

CD79a
Rabbit Monoclonal antibody(Mab)
Catalog # AD80015

Specification

CD79a - Product info

Application	IHC-P, IHC
Primary Accession	P11912
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	25038

CD79a - Additional info

Gene ID	973
Gene Name	CD79A

Other Names

B-cell antigen receptor complex-associated protein alpha chain, Ig-alpha, MB-1 membrane glycoprotein, Membrane-bound immunoglobulin-associated protein, Surface IgM-associated protein, CD79a, CD79A, IGA, MB1

Dilution

IHC-P~~Ready-to-use
IHC~~Ready-to-use

Storage	This product is stored at 2-8 °C, please use it within the expiration date.
Precautions	CD79a Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CD79a - Protein Information

Name CD79A

Synonyms
Function

IGA, MB1
Required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells. Stimulates SYK autophosphorylation and activation. Binds to BLNK, bringing BLNK into proximity with SYK and allowing SYK

Cellular Location

to phosphorylate BLNK. Also interacts with and increases activity of some Src- family tyrosine kinases. Represses BCR signaling during development of immature B-cells. Cell membrane; Single-pass type I membrane protein. Note=Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts. B-cells.


Tissue Location

CD79a - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD79a - Images



Tonsil



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using AD80015 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room

temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.