

TOP2A
Rabbit Monoclonal antibody(Mab)
Catalog # AD80039**Specification**

TOP2A - Product info

Application	IHC-P, IHC
Primary Accession	P11388
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	174385

TOP2A - Additional info

Gene ID	7153
Gene Name	TOP2A
Other Names	
DNA topoisomerase 2-alpha, 5.6.2.2 {ECO:0000255 PROSITE-ProRule:PRU00995, ECO:0000269 PubMed:19222228, ECO:0000269 PubMed:19697956}, DNA topoisomerase II, alpha isozyme, TOP2A, TOP2	

Dilution

IHC-P~~Ready-to-use
IHC~~Ready-to-use

Storage

Maintain refrigerated at 2-8°C

Precautions

TOP2A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TOP2A - Protein Information**Name TOP2A**

Synonyms
Function

TOP2
Control of topological states of DNA by transient breakage and subsequent rejoining of DNA strands. Topoisomerase II makes double-strand breaks. Essential during mitosis and meiosis for proper segregation of daughter chromosomes. May play a role in regulating the period length of ARNTL/BMAL1 transcriptional oscillation (By similarity). Cytoplasm. Nucleus, nucleoplasm. Note=Generally located in the nucleoplasm


Cellular Location

TOP2A - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TOP2A - Images



Tonsil



Immunohistochemical analysis of paraffin-embedded tissue using AD80294 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.