

Vimentin
Mouse Monoclonal antibody(Mab)
Catalog # AD80040

Specification

Vimentin - Product info

Application	IHC-P, IHC
Primary Accession	P08670
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	53652

Vimentin - Additional info

Gene ID	7431
Gene Name	VIM
Other Names	
Vimentin, VIM	

Dilution

IHC-P~~Ready-to-use
IHC~~Ready-to-use

Storage	This product is stored at 2-8 °C, please use it within the expiration date.
Precautions	Vimentin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Vimentin - Protein Information

Name VIM ([HGNC:12692](#))


Function	Vimentins are class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells. Vimentin is attached to the nucleus, endoplasmic reticulum, and mitochondria, either laterally or terminally.
Cellular Location	Cytoplasm. Cytoplasm, cytoskeleton. Nucleus matrix {ECO:0000250 UniProtKB:P31000}
Tissue Location	Highly expressed in fibroblasts, some expression in T- and B-lymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone- independent mammary carcinoma cell lines

Vimentin - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Vimentin - Images



Liver



Immunohistochemical analysis of paraffin-embedded gastrointestinal stromal tumor tissue using AD80305 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.