

**CD13**  
**Rabbit Monoclonal antibody(Mab)**  
**Catalog # AD80057**

## Specification

---

### CD13 - Product info

Application	<b>IHC-P, IHC</b>
Primary Accession	<a href="#">P15144</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal</b>
Calculated MW	<b>109540</b>

### CD13 - Additional info

Gene ID	<b>290</b>
Gene Name	<b>ANPEP</b>

#### Other Names

Aminopeptidase N, AP-N, hAPN, 3.4.11.2, Alanyl aminopeptidase, Aminopeptidase M, AP-M, Microsomal aminopeptidase, Myeloid plasma membrane glycoprotein CD13, gp150, CD13, ANPEP, APN, CD13, PEPN

#### Dilution

IHC-P~~Ready-to-use  
IHC~~Ready-to-use

Storage	<b>This product is stored at 2-8 °C, please use it within the expiration date.</b>
Precautions	<b>CD13 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.</b>

### CD13 - Protein Information

**Name ANPEP**

Synonyms  
Function

**APN, CD13, PEPN**

**Broad specificity aminopeptidase which plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Also involved in the processing of various peptides including peptide hormones, such as angiotensin III and IV, neuropeptides, and chemokines. May also be involved the cleavage of peptides bound to major histocompatibility complex class II molecules of antigen presenting cells. May have a role in angiogenesis and**

Cellular Location

**promote cholesterol crystallization. Cell membrane; Single-pass type II membrane protein. Note=Also found as a soluble form**

Tissue Location


**Expressed in epithelial cells of the kidney, intestine, and respiratory tract; granulocytes, monocytes, fibroblasts, endothelial cells, cerebral pericytes at the blood- brain barrier, synaptic membranes of cells in the CNS. Also expressed in endometrial stromal cells, but not in the endometrial glandular cells. Found in the vasculature of tissues that undergo angiogenesis and in malignant gliomas and lymph node metastases from multiple tumor types but not in blood vessels of normal tissues. A soluble form has been found in plasma. It is found to be elevated in plasma and effusions of cancer patients**

### CD13 - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CD13 - Images



Tonsil



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using AD80057 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.