

**IgG**  
**Rabbit Polyclonal antibody(Pab)**  
**Catalog # AD80082**

## Specification

---

### IgG - Product info

Application	IHC-P, IHC
Primary Accession	<a href="#">P01857</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43912

### IgG - Additional info

Gene Name	IGHG1 {ECO:0000303 PubMed:11340299, ECO:0000303 Ref.11}
-----------	---

#### Other Names

Immunoglobulin heavy constant gamma 1 {ECO:0000303|PubMed:11340299, ECO:0000303|Ref.13}, Ig gamma-1 chain C region, Ig gamma-1 chain C region EU, Ig gamma-1 chain C region KOL, Ig gamma-1 chain C region NIE, IGHG1 {ECO:0000303|PubMed:11340299, ECO:0000303|Ref.13}

#### Dilution

IHC-P~~Ready-to-use  
IHC~~Ready-to-use

#### Storage

#### Precautions

**This product is stored at 2-8 °C, please use it within the expiration date.**

**IgG Antibody is for research use only and not for use in diagnostic or therapeutic procedures.**

### IgG - Protein Information

**Name** IGHG1 {ECO:0000303|PubMed:11340299, ECO:0000303|Ref.13}

#### Function

**Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the**

effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:[22158414](#), PubMed:[20176268](#)). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:[17576170](#), PubMed:[20176268](#)).  
Secreted. Cell membrane


Cellular Location

### IgG - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### IgG - Images



Appendix



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using AD80082 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH 6.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.