

Lambda Chain
Mouse Monoclonal antibody(Mab)
Catalog # AD80089**Specification**

Lambda Chain - Product info

Application	IHC-P, IHC
Primary Accession	P01701
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	12249

Lambda Chain - Additional info

Gene Name **IGLC1 {ECO:0000303|PubMed:11872955, ECO:0000303|Ref.6}**

Other Names

Immunoglobulin lambda variable 1-51 {ECO:0000303|PubMed:11872955, ECO:0000303|Ref.7}, Ig lambda chain V-I region BL2, Ig lambda chain V-I region EPS, Ig lambda chain V-I region NEW, Ig lambda chain V-I region NIG-64, IGLV1-51 {ECO:0000303|PubMed:11872955, ECO:0000303|Ref.7}

Dilution

IHC-P~~Ready-to-use

IHC~~Ready-to-use

Storage

This product is stored at 2-8 °C, please use it within the expiration date.

Precautions

Lambda Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Lambda Chain - Protein Information

Name IGLV1-51 {ECO:0000303|PubMed:11872955, ECO:0000303|Ref.7}

Function

Constant region of immunoglobulin light chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which

results in the elimination of bound antigens (PubMed:[22158414](#), PubMed:[20176268](#)). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:[17576170](#), PubMed:[20176268](#)).
Secreted. Cell membrane


Cellular Location

Lambda Chain - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Lambda Chain - Images



Tonsil



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using AD80089 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH 6.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.