

Myosin Light Chain2
Rabbit Monoclonal antibody(Mab)
Catalog # AD80100**Specification**

Myosin Light Chain2 - Product info

Application	IHC-P, IHC
Primary Accession	P10916
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	18789

Myosin Light Chain2 - Additional info

Gene ID	4633
Gene Name	MYL2 (HGNC:7583)

Other Names

Myosin regulatory light chain 2, ventricular/cardiac muscle isoform, MLC-2, Ventricular myosin light chain 2, MYL2 (HGNC:7583)

Dilution

IHC-P~~Ready-to-use

IHC~~Ready-to-use

Storage

This product is stored at 2-226 °C, please use it within the expiration date.

Precautions

Myosin Light Chain2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Myosin Light Chain2 - Protein Information

Name MYL2 ([HGNC:7583](#))

Function

Contractile protein that plays a role in heart development and function (By similarity). Following phosphorylation, plays a role in cross-bridge cycling kinetics and cardiac muscle contraction by increasing myosin lever arm stiffness and promoting myosin head diffusion; as a consequence of the increase in maximum contraction force and calcium sensitivity of contraction force. These events altogether slow down myosin kinetics and prolong duty cycle resulting in accumulated myosins being cooperatively recruited to

actin binding sites to sustain thin filament activation as a means to fine-tune myofilament calcium sensitivity to force (By similarity). During cardiogenesis plays an early role in cardiac contractility by promoting cardiac myofibril assembly (By similarity).

Cellular Location


Cytoplasm, myofibril, sarcomere, A band
{ECO:0000250|UniProtKB:P08733}

Myosin Light Chain2 - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Myosin Light Chain2 - Images



Normal breast tissues



Immunohistochemical analysis of paraffin-embedded human normal breast tissue using AD80100 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.