

AR
Rabbit Monoclonal antibody(Mab)
Catalog # AD80215

Specification

AR - Product info

Application	IHC-P, IHC
Primary Accession	P10275
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	99188

AR - Additional info

Gene ID	367
Gene Name	AR

Other Names

Androgen receptor, Dihydrotestosterone receptor, Nuclear receptor subfamily 3 group C member 4, AR, DHTR, NR3C4

Dilution

IHC-P~~Ready-to-use
IHC~~Ready-to-use

Storage

Maintain refrigerated at 2-8°C

Precautions

Androgen Receptor Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

AR - Protein Information

Name AR

Synonyms
Function

DHTR, NR3C4
Steroid hormone receptors are ligand-activated transcription factors that regulate eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues. Transcription factor activity is modulated by bound coactivator and corepressor proteins like ZBTB7A that recruits NCOR1 and NCOR2 to the androgen response elements/ARE on target genes, negatively regulating androgen receptor signaling and androgen-induced cell proliferation

Cellular Location

(PubMed:[20812024](#)). Transcription activation is also down-regulated by NR0B2. Activated, but not phosphorylated, by HIPK3 and ZIPK/DAPK3.

Nucleus. Cytoplasm. Note=Detected at the promoter of target genes

(PubMed:[25091737](#)). Predominantly

cytoplasmic in unligated form but translocates to the nucleus upon ligand-binding. Can also translocate to the nucleus in unligated form in the presence of RACK1.

Tissue Location

Isoform 2 is mainly expressed in heart and skeletal muscle (PubMed:[15634333](#)).


Isoform 3 is expressed by basal and stromal cells of prostate (at protein level) (PubMed:[19244107](#)).

AR - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AR - Images



Prostate cancer



Immunohistochemical analysis of paraffin-embedded prostatic cancer tissue using AD80215 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.