



#### **SOX-10**

Rabbit Monoclonal antibody(Mab)
Catalog # AD80247

## **Specification**

### **SOX-10 - Product info**

Application IHC-P, IHC
Primary Accession P56693
Reactivity Human
Host Rabbit
Clonality Monoclonal
Calculated MW 49911

#### SOX-10 - Additional info

Gene ID 6663
Gene Name SOX10
Other Names

Dilution

Transcription factor SOX-10, SOX10

IHC-P~~Ready-to-use IHC~~Ready-to-use

**Storage**Maintain refrigerated at 2-8°C

Precautions SOX-10 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

# **SOX-10 - Protein Information**

Name SOX10

Function Transcription factor that plays a central

role in developing and mature glia.

Specifically activates expression of myelin genes, during oligodendrocyte (OL) maturation, such as DUSP15 and MYRF, thereby playing a central role in oligodendrocyte maturation and CNS myelination. Once induced, MYRF

cooperates with SOX10 to implement the myelination program. Transcriptional activator of MITF, acting synergistically

with PAX3 (PubMed: 21965087).

Cytoplasm. Nucleus. Mitochondrion outer

membrane

Cellular Location



Tel: 858.875.1900 Fax: 858.875.1999

{ECO:0000250|UniProtKB:Q04888}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q04888}; Cytoplasmic side {ECO:0000250|UniProtKB:Q04888} Expressed in fetal brain and in adult brain, heart, small intestine and colon

Tissue Location

### SOX-10 - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# SOX-10 - Images



## Malignant melanoma



Immunohistochemical analysis of paraffin-embedded 0 tissue using AD80278 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(Ready-to-use) for 15 min at room temperature. AmpSeeTM Detection Systems∏Abcepta:AR005∏ was used as the secondary antibody.