

p16
Mouse Monoclonal antibody(Mab)
Catalog # AD80303

Specification

p16 - Product info

| | |
|-------------------|------------------------|
| Application | IHC-P, IHC |
| Primary Accession | P42771 |
| Reactivity | Human |
| Host | Mouse |
| Clonality | Monoclonal |
| Calculated MW | 16533 |

p16 - Additional info

| | |
|-----------|--------------------------------------|
| Gene ID | 1029 |
| Gene Name | CDKN2A (HGNC:1787) |

Other Names

Cyclin-dependent kinase inhibitor 2A {ECO:0000312|HGNC:HGNC:1787}, Cyclin-dependent kinase 4 inhibitor A, CDK4I, Multiple tumor suppressor 1, MTS-1, p16-INK4a, p16-INK4, p16INK4A, CDKN2A (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=1787), CDKN2, MTS1

Dilution

IHC-P~~Ready-to-use
IHC~~Ready-to-use

Storage

Maintain refrigerated at 2-8°C

Precautions

p16 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

p16 - Protein Information

Name CDKN2A ([HGNC:1787](#))

Synonyms
Function

CDKN2, MTS1

Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.

Cellular Location
Tissue Location


Cytoplasm. Nucleus
Widely expressed but not detected in brain or skeletal muscle. Isoform 3 is pancreas-specific

p16 - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

p16 - Images



Cervical cancer



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma tissue using AD80303 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems [Abcepta:AR005] was used as the secondary antibody.