



#### **VHL**

Mouse Monoclonal antibody(Mab)
Catalog # AD80335

## **Specification**

### **VHL - Product info**

Application IHC-P, IHC
Primary Accession P40337
Reactivity Human
Host Mouse
Clonality Monoclonal
Calculated MW 24153

#### **VHL - Additional info**

Gene ID 7428
Gene Name VHL

**Other Names** 

von Hippel-Lindau disease tumor suppressor, Protein G7, pVHL, VHL

**Dilution** 

IHC-P~~Ready-to-use IHC~~Ready-to-use

Storage

Maintain refrigerated at 2-8°C

Precautions VHL Antibody is for research use only and

not for use in diagnostic or therapeutic

procedures.

#### **VHL - Protein Information**

**Name VHL** 

Function Involved in the ubiquitination and

subsequent proteasomal degradation via the von Hippel-Lindau ubiquitination complex. Seems to act as a target recruitment subunit in the E3 ubiquitin ligase complex and recruits hydroxylated hypoxia- inducible factor (HIF) under normoxic conditions. Involved in transcriptional repression through

interaction with HIF1A, HIF1AN and histone

deacetylases. Ubiquitinates, in an oxygen-responsive manner, ADRB2. Isoform 1: Cytoplasm. Membrane; Peripheral membrane protein. Nucleus.

Cellular Location



Tissue Location


Note=Found predominantly in the cytoplasm and with less amounts nuclear or membrane-associated Colocalizes with ADRB2 at the cell membrane Expressed in the adult and fetal brain and kidney

#### **VHL - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **VHL** - Images



## Colon cancer



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma tissue using AD80303 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(Ready-to-use) for 15 min at room temperature. AmpSeeTM Detection Systems Abcepta: AR005 was used as the secondary antibody.