

Myogenin
Rabbit Monoclonal antibody(Mab)
Catalog # AD80388

Specification

Myogenin - Product info

Application	IHC-P
Primary Accession	P15173
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	25037

Myogenin - Additional info

Gene ID	4656
---------	------

Other Names
Myogenin, Class C basic helix-loop-helix protein 3, bHLHC3, Myogenic factor 4, Myf-4, MYOG, BHLHC3, MYF4

Dilution
IHC-P~~Ready-to-use

Storage	This product is stored at 2-8 °C, please use it within the expiration date.
---------	--

Myogenin - Protein Information

Name MYOG

Synonyms	BHLHC3, MYF4
Function	Acts as a transcriptional activator that promotes transcription of muscle-specific target genes and plays a role in muscle differentiation, cell cycle exit and muscle atrophy. Essential for the development of functional embryonic skeletal fiber muscle differentiation. However is dispensable for postnatal skeletal muscle growth; phosphorylation by CAMK2G inhibits its transcriptional activity in response to muscle activity. Required for the recruitment of the FACT complex to muscle-specific promoter regions, thus promoting gene expression initiation. During terminal myoblast differentiation, plays a role as a strong activator of transcription at loci with an open chromatin structure previously initiated by

MYOD1. Together with MYF5 and MYOD1, co-occupies muscle-specific gene promoter core regions during myogenesis. Cooperates also with myocyte-specific enhancer factor MEF2D and BRG1-dependent recruitment of SWI/SNF chromatin-remodeling enzymes to alter chromatin structure at myogenic late gene promoters. Facilitates cell cycle exit during terminal muscle differentiation through the up-regulation of miR-20a expression, which in turn represses genes involved in cell cycle progression. Binds to the E-box containing (E1) promoter region of the miR-20a gene. Plays also a role in preventing reversal of muscle cell differentiation. Contributes to the atrophy-related gene expression in adult denervated muscles. Induces fibroblasts to differentiate into myoblasts (By similarity).

Cellular Location


Nucleus. Note=Recruited to late myogenic gene promoter regulatory sequences with SMARCA4/BRG1/BAF190A and SWI/SNF chromatin-remodeling enzymes to promote chromatin-remodeling and transcription initiation in developing embryos.

Myogenin - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Myogenin - Images



Alveolar rhabdomyosarcoma