

CTLA-4
Rabbit Monoclonal antibody(Mab)
Catalog # AD80586

Specification

CTLA-4 - Product info

Application	IHC-P
Primary Accession	P16410
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal
Calculated MW	24656

CTLA-4 - Additional info

Gene ID	1493
Other Names	
Cytotoxic T-lymphocyte protein 4, Cytotoxic T-lymphocyte-associated antigen 4, CTLA-4, CD152, CTLA4, CD152	
Storage	This product is stored at 2-8 °C, please use it within the expiration date.

CTLA-4 - Protein Information

Name CTLA4

Synonyms	CD152
Function	Inhibitory receptor acting as a major negative regulator of T-cell responses. The affinity of CTLA4 for its natural B7 family ligands, CD80 and CD86, is considerably stronger than the affinity of their cognate stimulatory coreceptor CD28.
Cellular Location	Cell membrane; Single-pass type I membrane protein. Note=Exists primarily an intracellular antigen whose surface expression is tightly regulated by restricted trafficking to the cell surface and rapid internalization
Tissue Location	Widely expressed with highest levels in lymphoid tissues. Detected in activated T-cells where expression levels are 30- to 50-fold less than CD28, the stimulatory coreceptor, on the cell surface following activation.

CTLA-4 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CTLA-4 - Images