

Functional Rab6-GTP Antibody, mAb (recombinant) (ATTO488)
Catalog # ADP0035

Specification

Functional Rab6-GTP Antibody, mAb (recombinant) (ATTO488) - Product Information

Application	ICC
Primary Accession	P20340
Reactivity	Human, Mouse, Drosophila
Host	Purified From HEK 293 Cell culture Supernatant.
Clonality	Monoclonal
Isotype	Human IgG2 λ
Gene Source	Human
Application Note	ICC(1:1'000)
Calculated MW	23593
Description	<p>anti-Rab6-GTP monoclonal antibody (recombinant) (AA2) is composed of human variable regions (VH and VL) (λ-chain) of immunoglobulin fused to the human IgG2 Fc domain.</p> <p>anti-Rab6-GTP monoclonal antibody (recombinant) (AA2) is an antibody developed by antibody phage display technology using a human naive antibody gene library. These libraries consist of scFv (single chain fragment variable) composed of VH (variable domain of the human immunoglobulin heavy chain) and VL (variable domain of the human immunoglobulin light chain) connected by a polypeptide linker. The antibody fragments are displayed on the surface of filamentous bacteriophage (M13). This scFv was selected by affinity selection on antigen in a process termed panning. Multiple rounds of panning are performed to enrich for antigen-specific scFv-phage. Monoclonal antibodies are subsequently identified by screening after each round of selection. The selected monoclonal scFv is cloned into an appropriate vector containing a Fc portion of interest and then produced in mammalian cells to generate an IgG like scFv-Fc fusion protein.</p>

Functional Rab6-GTP Antibody, mAb (recombinant) (ATTO488) - Additional Information

Gene ID 5870

Other Names

Ras-related Protein Rab-6

Target/Specificity

Recognizes human, mouse and Drosophila GTP-bound Rab6a and Rab6b and mutant Rab6Q72L. Does not detect Rab6•GDP.

Format

Liquid. In PBS containing 10% glycerol and 0.02% sodium azide.

Reconstitution & Storage

Stable for at least 1 month after receipt when stored at +4°C. Stable for at least 1 year after receipt when stored at -20°C.

Precautions

Functional Rab6-GTP Antibody, mAb (recombinant) (ATTO488) is for research use only and not for use in diagnostic or therapeutic procedures.

Functional Rab6-GTP Antibody, mAb (recombinant) (ATTO488) - Protein Information

Name RAB6A

Synonyms RAB6

Function

Regulator of COPI-independent retrograde transport from the Golgi apparatus towards the endoplasmic reticulum (ER) (PubMed:25962623). Has a low GTPase activity (PubMed:25962623). Recruits VPS13B to the Golgi membrane (PubMed:25492866). Plays a role in neuron projection development (Probable).

Cellular Location

Golgi apparatus membrane; Lipid- anchor. Cytoplasmic vesicle, secretory vesicle, acrosome membrane {ECO:0000250|UniProtKB:P35279}; Peripheral membrane protein. Note=BICD2 facilitates its targeting to Golgi apparatus membrane. [Isoform 2]: Golgi apparatus membrane; Lipid-anchor

Tissue Location

Ubiquitous..


Functional Rab6-GTP Antibody, mAb (recombinant) (ATTO488) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

Functional Rab6-GTP Antibody, mAb (recombinant) (ATTO488) - Images



Rab6-GTP is detected by immunocytochemistry using anti-Rab6-GTP, mAb (AA2).

Method: HeLa cells are grown in standard culture conditions, fixed with paraformaldehyde (3%), permeabilized in PBS+ BSA 0.2 % + Saponin 0.05 % and incubated with anti-Rab6-GTP, mAb (AA2) (1ug /ml) in PBS-BSA-Saponin. After incubation for 30 min at RT and several washes in PBS, cells are treated with a goat anti-human (Cy3) antibody in PBS-BSA-Saponin for 30 min at RT, washed and mounted in Moewiol. Nuclei are stained with DAPI.

Picture courtesy of Dr Moutel, Dr Franck Perez Lab, Curie Institute, Paris.

Functional Rab6-GTP Antibody, mAb (recombinant) (ATTO488) - Background

Rab6 is involved in protein transport. It is a regulator of membrane traffic from the Golgi apparatus towards the endoplasmic reticulum (ER).