

**Goat Anti-BIM Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF1154a

**Specification**

---

**Goat Anti-BIM Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">O43521</a>
Other Accession	<a href="#">NP_619529</a> , <a href="#">10018</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	22171

**Goat Anti-BIM Antibody - Additional Information**

**Gene ID** 10018

**Other Names**

Bcl-2-like protein 11, Bcl2-L-11, Bcl2-interacting mediator of cell death, BCL2L11, BIM

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-BIM Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-BIM Antibody - Protein Information**

**Name** BCL2L11

**Synonyms** BIM

**Function**

Induces apoptosis and anoikis. Isoform BimL is more potent than isoform BimEL. Isoform Bim-alpha1, isoform Bim-alpha2 and isoform Bim-alpha3 induce apoptosis, although less potent than isoform BimEL, isoform BimL and isoform BimS. Isoform Bim-gamma induces apoptosis. Isoform Bim-alpha3 induces apoptosis possibly through a caspase- mediated pathway. Isoform BimAC and isoform BimABC lack the ability to induce apoptosis.

### Cellular Location

Endomembrane system; Peripheral membrane protein. Note=Associated with intracytoplasmic membranes. [Isoform BimL]: Mitochondrion. [Isoform Bim-alpha1]: Mitochondrion.

### Tissue Location

Isoform BimEL, isoform BimL and isoform BimS are the predominant isoforms and are widely expressed with tissue-specific variation. Isoform Bim-gamma is most abundantly expressed in small intestine and colon, and in lower levels in spleen, prostate, testis, heart, liver and kidney.

### Goat Anti-BIM Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Goat Anti-BIM Antibody - Images



AF1154a staining (0.5  $\mu$ g/ml) of K562 lysate (RIPA buffer, 35  $\mu$ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.