

Goat Anti-BNIP1 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1162a**Specification**

Goat Anti-BNIP1 Antibody - Product Information

Application	WB
Primary Accession	Q12981
Other Accession	NP_053583 , 662 , 224630 (mouse) , 140932 (rat)
Reactivity	Human
Predicted	Mouse, Rat
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	26132

Goat Anti-BNIP1 Antibody - Additional Information**Gene ID** 662**Other Names**

Vesicle transport protein SEC20, BCL2/adenovirus E1B 19 kDa protein-interacting protein 1, Transformation-related gene 8 protein, TRG-8, BNIP1, NIP1, SEC20L

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-BNIP1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-BNIP1 Antibody - Protein Information**Name** BNIP1**Synonyms** NIP1, SEC20L**Function**

As part of a SNARE complex may be involved in endoplasmic reticulum membranes fusion and be required for the maintenance of endoplasmic reticulum organization (PubMed:15272311). Also plays a

role in apoptosis (PubMed:15272311, PubMed:23896122, PubMed:7954800). It is for instance required for endoplasmic reticulum stress-induced apoptosis (PubMed:23896122). As a substrate of RNF185 interacting with SQSTM1, might also be involved in mitochondrial autophagy (Probable).

Cellular Location

Endoplasmic reticulum membrane; Single-pass type IV membrane protein. Mitochondrion membrane; Single-pass type IV membrane protein. Note=Localization to the mitochondrion is regulated by RNF186.

Tissue Location

Isoform 1 is highly expressed in heart, brain, liver skeletal muscle and pancreas. Isoform 3 is moderately expressed in placenta, lung and kidney. Isoform 4 is highly expressed in testis and small intestine.

Goat Anti-BNIP1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-BNIP1 Antibody - Images



AF1162a (0.5 µg/ml) staining of Human Kidney lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-BNIP1 Antibody - Background

This gene is a member of the BCL2/adenovirus E1B 19 kd-interacting protein (BNIP) family. It interacts with the E1B 19 kDa protein which is responsible for the protection of virally-induced cell death, as well as E1B 19 kDa-like sequences of BCL2, also an apoptotic protector. Alternative

splicing of this gene results in four protein products with identical N- and C-termini.

Goat Anti-BNIP1 Antibody - References

Polymorphisms in mitochondrial genes and prostate cancer risk. Wang L, et al. Cancer Epidemiol Biomarkers Prev, 2008 Dec. PMID 19064571.
The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.
Involvement of BNIP1 in apoptosis and endoplasmic reticulum membrane fusion. Nakajima K, et al. EMBO J, 2004 Aug 18. PMID 15272311.
Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932.
Novel BNIP1 variants and their interaction with BCL2 family members. Zhang H, et al. FEBS Lett, 1999 Apr 1. PMID 10217402.