

Goat Anti-DKK1 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1323a

Specification

Goat Anti-DKK1 Antibody - Product Information

| | |
|-------------------|---------------------------------------------------|
| Application | WB |
| Primary Accession | O94907 |
| Other Accession | NP_036374 , 22943 |
| Reactivity | Human |
| Predicted | Rat, Pig, Dog |
| Host | Goat |
| Clonality | Polyclonal |
| Concentration | 0.5 mg/ml |
| Isotype | IgG |
| Calculated MW | 28672 |

Goat Anti-DKK1 Antibody - Additional Information

Gene ID 22943

Other Names

Dickkopf-related protein 1, Dickkopf-1, Dkk-1, hDkk-1, SK, DKK1

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-DKK1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-DKK1 Antibody - Protein Information

Name DKK1

Function

Antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6 (PubMed:22000856). DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and

Alzheimer disease (PubMed:17143291). Inhibits the pro-apoptotic function of KREMEN1 in a Wnt-independent manner, and has anti-apoptotic activity (By similarity).

Cellular Location

Secreted.

Tissue Location

Placenta.

Goat Anti-DKK1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-DKK1 Antibody - Images



AF1323a (1 µg/ml) staining of human bone marrow lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-DKK1 Antibody - Background

This gene encodes a protein that is a member of the dickkopf family. It is a secreted protein with two cysteine rich regions and is involved in embryonic development through its inhibition of the WNT signaling pathway. Elevated levels of DKK1 in bone marrow plasma and peripheral blood is associated with the presence of osteolytic bone lesions in patients with multiple myeloma.

Goat Anti-DKK1 Antibody - References

Analysis of copy number variation in 8,842 Korean individuals reveals 39 genes associated with hepatic biomarkers AST and ALT. Kim HY, et al. BMB Rep, 2010 Aug. PMID 20797317. Maternal genes and facial clefts in offspring: a comprehensive search for genetic associations in two

population-based cleft studies from Scandinavia. Jugessur A, et al. PLoS One, 2010 Jul 9. PMID 20634891. Evaluation of candidate stromal epithelial cross-talk genes identifies association between risk of serous ovarian cancer and TERT, a cancer susceptibility hot-spot. Johnatty SE, et al. PLoS Genet, 2010 Jul 8. PMID 20628624. Dkk1 stabilizes Wnt co-receptor LRP6: implication for Wnt ligand-induced LRP6 down-regulation. Li Y, et al. PLoS One, 2010 Jun 8. PMID 20543981. Genetic association study of KREMEN1 and DKK1 and schizophrenia in a Japanese population. Aleksic B, et al. Schizophr Res, 2010 May. PMID 20153141.