

**Goat Anti-GAPDH (C Terminus) Loading Control Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF1464a

### Specification

---

### Goat Anti-GAPDH (C Terminus) Loading Control Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P04406</a>
Other Accession	<a href="#">NP_002037</a> , <a href="#">2597</a>
Reactivity	Human, Mouse, Rat, Pig
Predicted	Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5mg/ml
Isotype	IgG
Calculated MW	36053

### Goat Anti-GAPDH (C Terminus) Loading Control Antibody - Additional Information

Gene ID 2597

#### Other Names

Glyceraldehyde-3-phosphate dehydrogenase, GAPDH, 1.2.1.12, Peptidyl-cysteine S-nitrosylase  
GAPDH, 2.6.99.-, GAPDH, GAPD

#### Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

Goat Anti-GAPDH (C Terminus) Loading Control Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### Goat Anti-GAPDH (C Terminus) Loading Control Antibody - Protein Information

**Name** GAPDH {ECO:0000303|PubMed:2987855, ECO:0000312|HGNC:HGNC:4141}

#### Function

Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively (PubMed:<a href="http://www.uniprot.org/citations/11724794" target="\_blank">11724794</a>, PubMed:<a href="http://www.uniprot.org/citations/3170585" target="\_blank">3170585</a>). Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first

step of the pathway by converting D- glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate (PubMed:<a href="http://www.uniprot.org/citations/11724794" target="\_blank">11724794</a>, PubMed:<a href="http://www.uniprot.org/citations/3170585" target="\_blank">3170585</a>). Modulates the organization and assembly of the cytoskeleton (By similarity). Facilitates the CHP1- dependent microtubule and membrane associations through its ability to stimulate the binding of CHP1 to microtubules (By similarity). Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes (PubMed:<a href="http://www.uniprot.org/citations/23071094" target="\_blank">23071094</a>). Upon interferon-gamma treatment assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation (PubMed:<a href="http://www.uniprot.org/citations/23071094" target="\_blank">23071094</a>). Also plays a role in innate immunity by promoting TNF-induced NF-kappa-B activation and type I interferon production, via interaction with TRAF2 and TRAF3, respectively (PubMed:<a href="http://www.uniprot.org/citations/23332158" target="\_blank">23332158</a>, PubMed:<a href="http://www.uniprot.org/citations/27387501" target="\_blank">27387501</a>). Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis (By similarity). Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC (By similarity).

#### Cellular Location

Cytoplasm, cytosol. Nucleus {ECO:0000250|UniProtKB:P04797}. Cytoplasm, perinuclear region. Membrane Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P04797} Note=Translocates to the nucleus following S-nitrosylation and interaction with SIAH1, which contains a nuclear localization signal (By similarity). Postnuclear and Perinuclear regions (PubMed:12829261) {ECO:0000250|UniProtKB:P04797, ECO:0000269|PubMed:12829261}


#### Goat Anti-GAPDH (C Terminus) Loading Control Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Goat Anti-GAPDH (C Terminus) Loading Control Antibody - Images






AF1464a (0.1 µg/ml) staining of lysates of cell line HEK293 (A) and HeLa (B) (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF1464a (0.1ug/ml) staining of PFA-fixed HeLa before (top) and after (bottom) si-RNA-mediated GAPDH knock-down expression. Primary incubation 1h at ambient temp. Detection by DyLight 488. Nuclear DAPI stain (right). This data was obtained using a previous



AF1464a (0.5 µg/ml) staining of Mouse Liver (A) and Rat Brain (B) lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

### **Goat Anti-GAPDH (C Terminus) Loading Control Antibody - Background**

The product of this gene catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The enzyme exists as a tetramer of identical chains. Many pseudogenes similar to this locus are present in the human genome.

### **Goat Anti-GAPDH (C Terminus) Loading Control Antibody - References**

Inhibition of glyceraldehyde-3-phosphate dehydrogenase activity by antibodies present in the cerebrospinal fluid of patients with multiple sclerosis. Kϫlln J, et al. J Immunol, 2010 Aug 1. PMID 20610654.

Proteome analysis of the thalamus and cerebrospinal fluid reveals glycolysis dysfunction and potential biomarkers candidates for schizophrenia. Martins-de-Souza D, et al. J Psychiatr Res, 2010 May 14. PMID 20471030.

Sex-specific proteome differences in the anterior cingulate cortex of schizophrenia.

Martins-de-Souza D, et al. J Psychiatr Res, 2010 Apr 8. PMID 20381070.

Identification of melanoma antigens using a Serological Proteome Approach (SERPA). Suzuki A, et al. Cancer Genomics Proteomics, 2010 Jan-Feb. PMID 20181627.

siah-1 Protein is necessary for high glucose-induced glyceraldehyde-3-phosphate dehydrogenase nuclear accumulation and cell death in Muller cells. Yego EC, et al. J Biol Chem, 2010 Jan 29. PMID 19940145.